



California State Board of Pharmacy
2720 Gateway Oaks Drive, Suite 100
Sacramento, CA 95833
Phone: (916) 518-3100 Fax: (916) 574-8618
www.pharmacy.ca.gov

Business, Consumer Services and Housing Agency
Department of Consumer Affairs
Gavin Newsom, Governor



LICENSING COMMITTEE REPORT

Debbie Veale, Licensee Member, Chairperson
Seung Oh, License Member, Vice-Chairperson
Lavanza Butler, Licensee Member
Jignesh Patel, Licensee Member
Jason Weisz, Public Member
Albert Wong, Licensee Member

The Licensing Committee will meet on January 27, 2021. An update of the work of the committee will be provided during the board meeting

a. Summary of Presentation by the University of California Schools of Pharmacy related to Academic Dishonesty

Background

As part of its July 2020 Board Meeting, the Board received a presentation on recently published research regarding academic dishonesty in the California Schools of Pharmacy. Subsequent to that presentation, the Licensing Committee as part of its October 2020 meeting, discussed the issue and possible actions the Board could take to address the issue. As the minutes reflect, no action was taken during the committee, but the committee indicated its desire to continue its discussion.

During the Meeting

During the committee meeting members will receive a summary of the presentation provided by representatives of the University of California, including its approach to academic dishonesty and best practices for creating an environment that discourages such behavior.

b. Discussion and Consideration of Proposal to Expand the Authority for Pharmacists to Order and Administer CLIA Waived Tests for Influenza and COVID

Relevant Law

Business and Professions Code (BPC) 4052(a)(12) establishes the authority for a pharmacist to order and interpret tests for the purpose of monitoring and managing the efficacy and toxicity of drug therapies. As included in this provision, the pharmacist performing such functions must ensure such testing is done in coordination with the patient's primary care provider or diagnosing prescriber, as appropriate.

BPC 4052.1(a)(2) establishes the authority for a pharmacist to order drug therapy-related laboratory tests in a licensed health care facility in accordance with policies, procedures, or protocols developed by health professionals, as specified.

BPC 4052.2(a)(2) establishes the authority for a pharmacist to order drug therapy-related laboratory tests as part of the care provided in a licensed health care facility, licensed home health agency, licensed correctional clinic, a licensed clinic with physician oversight, or other provider as specified, in accordance with the policies, procedures, or protocols of that facility, home health agency, etc.

BPC 4052.4 establishes the authority for a pharmacist to perform skin puncture in the course of performing routine patient assessment procedures or in the course of performing any procedure authorized under BPC 1206.5 or BPC 1206.6. The section provides that “routine patient assessment procedures,” includes CLIA waived tests as authorized under BPC 1206.5 and 1206.6.

BPC 1206.5(a)(11) establishes the authority for a pharmacist to perform a clinical laboratory test or examination classified as waived under CLIA as long as the clinical laboratory test or examination is performed under the overall operation and administration of the laboratory director, as described in Section 1209, including, but not limited to, documentation by the laboratory director of the adequacy of the qualifications and competency of the personnel.

BPC 1206.6 provides authority for pharmacist at a community pharmacy who, performs only blood glucose, hemoglobin A1c, or cholesterol tests that are classified as waived under CLIA in the course of performing assessments as provided in BPC 4052.4. This section also requires the pharmacy to obtain a CLIA certificate of waiver and comply with all other requirements for the performance of waived clinical laboratory tests under applicable federal regulations. Further, the section provides that the pharmacist-in-charge (PIC) is responsible for directing and supervising testing oversight, decision making and ensures the pharmacy has obtained a registration as required by BPC 1265.

BPC 1265 establishes the licensing requirements for a clinical laboratory as specified. BPC 1265(k) provides authority for the PIC to serve as the laboratory director for registration required under BPC 1206.6.

Background

On August 25, 2020, the DCA Director issued an order that waives specified professional licensing requirements and amends the scopes of practice of pharmacists and pharmacy technicians to allow them to perform waived, point-of-care tests used to detect SARS-CoV-2. Along with the waiver, guidance was released to inform and educate pharmacies, pharmacists and pharmacy technician of clinical laboratory requirements that apply under the DCA Order.

More recently, as part of the October Licensing Committee Meeting and subsequent Board Meeting, the Board approved the following policy statement:

The CDC has acknowledged that the flu and COVID-19 are both respiratory illnesses that are caused by different viruses that may be difficult to differentiate based on symptoms alone without testing to confirm a diagnosis. The Board also recognizes that community pharmacies provide unique access for patients to obtain tests in a safe and convenient location. In recognition of these facts and the existing authority pharmacists already may provide certain CLIA waived tests, the Board hereby declares its support for all efforts to secure temporary authority for pharmacists to perform CLIA-waived tests for influenza and COVID during the declared disaster, as well as a more permanent solution through statutory changes that facilitate authority for pharmacists to perform CLIA-waived COVID and influenza testing in a safe manner.

During the Meeting

During the meeting members will have the opportunity to consider a draft statutory proposal if approved and recommended by the Committee.

Attachment 1 includes the draft statutory proposal that will be considered by the Committee.

c. Discussion and Consideration of Statutory Proposal to Expand the Authority for Pharmacy Technicians to Administer COVID-19 and Influenza Vaccines

Relevant Law

BPC 4115 provides that a pharmacy technician may perform packaging, manipulative, repetitive, or other nondiscretionary tasks only while assisting, and while under the direct supervision and control of, a pharmacist. This section further requires the board to adopt regulations that a pharmacy technician may perform.

Title 16, California Code of Regulations section 1793.2 establishes the nondiscretionary tasks to include removing the drug or drugs from stock; counting, pouring, or mixing pharmaceuticals, placing the product into the container, affixing the label or labels to the container; and packaging or repackaging.

Background

Existing law provides authority for pharmacists to independently initiate and administer vaccines lists on the routine schedules recommended by the federal Advisory Committee on Immunization Practices (ACIP) under specific conditions (BPC 4052.8). Effective January 1, 2021, this authority was expanded to include COVID-19 vaccines that are FDA authorized or FDA approved.

As part of the October Committee meeting and subsequent discussions during the October and November 2020 Board meetings, the Board approved the following policy statement related to pharmacy technician administered vaccinations.

Policy Statement – Expand Authority to Allow Pharmacy Technicians to Administer COVID-19 and Influenza Vaccinations

In recognition of the current COVID-19 crisis and consistent with the recommendations from health experts, including the CDC, on the importance of influenza and COVID-19 vaccinations, the Board supports all efforts to facilitate influenza and COVID-19 administration in a safe manner. Further, in recognition of the unique access patients have to community pharmacies, such locations provide a safe and convenient option to receive such vaccinations. The Board further believes that influenza and COVID-19 vaccine administration may be safely performed by a specially trained licensed pharmacy technician under specified conditions and as such supports efforts to secure such temporary authority under waivers during the declared disaster, as well as a more permanent solution through statutory or regulatory changes.

During the Meeting

During the meeting, members will receive an update on the Committee’s efforts to finalize a draft statutory proposal. The proposed under consideration of the Committee includes prior provisions identified by the Committee and approved by the Board including:

- Specificity that the task must be delegated by the supervising pharmacist
- Completion of a training program approved by ACPE and CPR certification
- Ongoing requirement for 1 hour of CE
- Authority to administer epinephrine, if delegated by the supervising pharmacist.
- Recordkeeping requirements

Attachment 2 includes a copy of the relevant laws and draft statutory proposal that will be considered by the Committee.

d. Discussion and Consideration of Board’s Current Policy related to Authority for Pharmacy Technicians to Administer Vaccines to Determine if Inclusion of Additional Vaccines is Appropriate

Background

The history of prior policy discussion is provided under the prior agenda item.

During the Meeting

During the meeting members will have the opportunity to receive an updated from the Committee on its discussion regarding the current policy statement and any recommendations it offered to the Board regarding expansion of the policy to include all vaccines included on the ACIP recommended schedule.

Should the Committee determine that it is appropriate to expand the policy, it is recommended that the policy statement be updated to reflect this expansion and conforming changes made to the draft statutory proposal discussed as part of the prior agenda item.

Attachment 3 includes a copy of the Recommended Child and Adolescent Immunization Schedule and Recommended Adult Immunization Schedule.

e. Discussion and Consideration of Draft Pharmacist Workforce Survey

Background

As indicated in the Board's responses to Sunset Issues, the issue of medication errors must be addressed to improve patient health. The issue warrants study in California, where conditions within a pharmacy may be different than on a national level. Further, consideration should be given to determine if the Board or some other entity should receive reports of medication errors to gain a better understanding of the scope of the issue and report on the findings. It appears appropriate to conduct a survey on working conditions to ascertain if conditions in California may be a contributing factor.

During the meeting

During the meeting members will receive an update on the Committee's discussion of the draft workforce survey. The draft survey was developed in collaboration with an expert from the Department of Consumer Affairs.

Attachment 4 includes a copy of the draft survey that will be reviewed by the Committee as part of its meeting.

I. Discussion and Consideration of Waiver Request of Business and Professions Code Section 4131(b) Related to the Location of the Supervising Pharmacy and Remote Dispensing Site Pharmacy

Subsequent to the release of the agenda, the request for consideration was withdrawn. No action is required at this time.

II. Review and Discussion of Licensing Statistics

The quarterly licensing statistics for fiscal year 2020/2021, are provided in **Attachment 5**.

As of December 31, 2020, the Board has received 7,623 initial applications, including:

- 1,348 intern pharmacists
- 1,525 pharmacist exam applications (451 new, 1,074 retake)
- 104 advanced practice pharmacists
- 2,218 pharmacy technicians
- 211 community pharmacy license applications

- 62 sterile compounding pharmacy license applications (LSC, LSE, SCP, SCE)
- 63 nonresident pharmacy license applications
- 16 hospital pharmacy license applications

As of December 31, 2020, the Board has received 275 requests for temporary site license applications, including:

- 135 community pharmacy license applications
- 38 sterile compounding pharmacy license applications
- 39 nonresident pharmacy license applications
- 18 hospital pharmacy license applications

As of December 31, 2020, the Board has issued 4,525 individual licenses, including:

- 1,324 intern pharmacists
- 1,493 pharmacists
- 75 advanced practice pharmacists
- 1,473 pharmacy technicians

As of December 31, 2020, the Board has issued 241 site licenses without temporary license requests, including:

- 82 automated drug delivery systems
- 44 community pharmacies
- 0 hospital pharmacies

As of December 31, 2020, the Board has issued 193 temporary site licenses, including:

- 98 community pharmacies
- 11 hospital pharmacies

Processing Times

The general application and deficiency mail processing times by license type are provided below reflecting data current as of January 15, 2021. The data reflects the time from when an application or deficiency response is received by the Board through to the time it is reviewed by licensing staff. The standard performance processing time is within 30 days for initial applications and is within 10 days for deficiency mail. The term “Current” means there are no items to review or staff is currently reviewing the items within 1-5 days for that specific license type.

Staff continue to work diligently and make adjustments to review applications and mail electronically during this pandemic. Processing times exceed the standard processing times resulting from a combination of factors, including impacts resulting from the pandemic.

Premises Application Types	Application Processing Times as of 10/2/2020	Application Processing Times as of 1/15/2021	Deficiency Mail Processing Times as of 10/2/2020	Deficiency Mail Processing Times as of 1/15/2021
Pharmacy	28	47	10	47
Nonresident Pharmacy	14	58	10	65
Sterile Compounding	28	16	22	33
Nonresident Sterile Compounding	22	Current	21	38
Outsourcing	Current	Current	Current	Current
Nonresident Outsourcing	17	19	Current	25
Hospital Satellite Compounding Pharmacy	Current	Current	Current	Current
Hospital	2	16	Current	Current
Clinic	22	42	10	3
Wholesaler	23	15	Current	18
Nonresident Wholesaler	24	22	Current	25
Third-Party Logistics Provider	Current	Current	Current	Current
Nonresident Third-Party Logistics Provider	8	11	Current	22
Automated Drug Delivery System	8	7	Current	Current
Automated Patient Dispensing System	Current	Current	Current	Current
Emergency Medical Services Automated Drug Delivery System	Current	Current	Current	Current

Individual Application Type	Application Processing Times as of 10/2/2020	Application Processing Times as of 1/15/2021	Application Processing Times as of 10/2/2020	Deficiency Mail Processing Times as of 1/15/2021
Exam Pharmacist	24	30	7	18
Pharmacist Initial Licensure	Current	Current	n/a	n/a
Advanced Practice Pharmacist	Current	28	Current	28

Individual Application Type	Application Processing Times as of 10/2/2020	Application Processing Times as of 1/15/2021	Application Processing Times as of 10/2/2020	Deficiency Mail Processing Times as of 1/15/2021
Intern Pharmacist	42	37	9	28
Pharmacy Technician	51	54	Current	34
Designated Representative	Current	31	Current	25
Designated Representatives-3PL	Current	29	Current	25
Designated Representatives-Reverse Distributor	Current	Current	Current	Current
Designated Paramedic	Current	Current	Current	Current

h. Future Meeting Dates

- April 21, 2021
- July 14, 2021
- October 27, 2021

Attachment 1

Draft Statutory Proposal Related to Pharmacist Provided COVID-19 and Influenza Testing

Proposal to Amend Business and Professions Code section 4052.4.

(a) Notwithstanding Section 2038 or any other provision of law, a pharmacist may perform skin puncture in the course of performing routine patient assessment procedures or in the course of performing any procedure authorized under Section 1206.5 or 1206.6. For purposes of this section, "routine patient assessment procedures" means: (a) procedures that a patient could, with or without a prescription, perform for himself or herself, or (b) clinical laboratory tests that are classified as waived pursuant to the federal Clinical Laboratory Improvement Amendments of 1988 (42 U.S.C. Sec. 263a) and the regulations adopted thereunder by the federal Health Care Financing Administration, as authorized by paragraph (11) of subdivision (a) of Section 1206.5 or Section 1206.6. A pharmacist performing these functions shall report the results obtained from a test to the patient and any physician designated by the patient. Any pharmacist who performs the service authorized by this section shall not be in violation of Section 2052.

(b) A pharmacist may perform any aspect of any FDA approved or authorized point-of-care test for the presence of SARS-CoV-2 or influenza that are classified as waived pursuant to the federal Clinical Laboratory Improvement Amendments as described in (a) under the following conditions:

1. The pharmacist completes the testing in a pharmacy licensed by the Board and that is appropriately licensed in California as a laboratory pursuant to BPC 1265.
2. The pharmacist has completed necessary training as specified in the pharmacy's policies and procedures.

Add BPC section 4119.10

A pharmacy located in this state, may employ pharmacists to perform FDA approved or authorized point-of-care tests for the presence of SARS-CoV-2 or influenza that are classified as waived pursuant the Federal Clinical Laboratory Improvement Amendments of 1988 under the following conditions:

1. The pharmacy is appropriately licensed as a laboratory under BPC section 1265.
2. The pharmacy maintains policies and procedures that at minimum describe the following:
 - a. Establish the initial training requirements, including specimen collection techniques relevant to the test(s) being performed at the pharmacy and ongoing training.
 - b. Establish the necessary safety precautions to protect pharmacy staff and consumers to reduce the risk of transmission consistent with CalOSHA and CDC requirements. Such policies should, at a minimum, include provisions for use for personal protective equipment, cleaning and sanitizing procedures, appropriate biohazard waste requirements and space requirements to protect the safety of staff and consumers.
 - c. Ensure dedicated physical or other segregated space that allows for privacy during the testing process, provides for private consultation with the pharmacist and to limit the potential contamination of other consumers in the pharmacy.

- d. Detail requirements for providing test results to the patient in a nonverbal manner, complying with mandatory reporting requirements to local and state reporting systems, and notification to primary care providers if consent is provided.
 - e. Ensure documentation of testing equipment maintenance and calibration.
 - f. Ensure appropriate storage and handling of specimens, testing reagents, etc.
3. The pharmacist-in-charge must review the policies and procedures on an annual basis. As part of this annual review the pharmacist-in-charge must also assess the pharmacy's compliance with its policies and where noncompliance is noted, document corrective actions to be taken. Documentation of the review must be maintained in a readily retrievable format for a period of three years from the date of completion.
4. The pharmacy must maintain documentation related to performing these tests that demonstrate compliance with all conditions in this subsection, including, the name of the pharmacist performing the test, the results and communication of results to a patient's primary medical provider. These documents must be maintained for period of three years from the date of making and must be maintained in a readily retrievable format.

Amend BPC section 1206.5.

(a) Notwithstanding subdivision (b) of Section 1206 and except as otherwise provided in Sections 1206.6 and 1241, no person shall perform a clinical laboratory test or examination classified as waived under CLIA unless the clinical laboratory test or examination is performed under the overall operation and administration of the laboratory director, as described in Section 1209, including, but not limited to, documentation by the laboratory director of the adequacy of the qualifications and competency of the personnel, and the test is performed by any of the following persons:

- (1) A licensed physician and surgeon holding a M.D. or D.O. degree.
- (2) A licensed podiatrist, a licensed dentist, or a licensed naturopathic doctor, if the results of the tests can be lawfully utilized within his or her practice.
- (3) A person licensed under this chapter to engage in clinical laboratory practice or to direct a clinical laboratory.
- (4) A person authorized to perform tests pursuant to a certificate issued under Article 5 (commencing with Section 101150) of Chapter 2 of Part 3 of Division 101 of the Health and Safety Code.
- (5) A licensed physician assistant if authorized by a supervising physician and surgeon in accordance with Section 3502 or 3535.
- (6) A person licensed under Chapter 6 (commencing with Section 2700).
- (7) A person licensed under Chapter 6.5 (commencing with Section 2840).

- (8) A perfusionist if authorized by and performed in compliance with Section 2590.
 - (9) A respiratory care practitioner if authorized by and performed in compliance with Chapter 8.3 (commencing with Section 3700).
 - (10) A medical assistant, as defined in Section 2069, if the waived test is performed pursuant to a specific authorization meeting the requirements of Section 2069.
 - (11) A pharmacist, as defined in Section 4036, if ordering drug therapy-related laboratory tests in compliance with paragraph (2) of subdivision (a) of Section 4052.1 or paragraph (2) of subdivision (a) of Section 4052.2, ~~or~~ if performing skin puncture in the course of performing routine patient assessment procedures in compliance with Section 4052.1 or performing testing as authorized in section 4052.4.
 - (12) A naturopathic assistant, as defined in Sections 3613 and 3640.2, if the waived test is performed pursuant to a specific authorization meeting the requirements of Sections 3613 and 3640.2.
 - (13) A licensed optometrist as authorized under Chapter 7 (commencing with Section 3000).
 - (14) Other health care personnel providing direct patient care.
 - (15) Any other person performing nondiagnostic testing pursuant to Section 1244.
- (b) Notwithstanding subdivision (b) of Section 1206, no person shall perform clinical laboratory tests or examinations classified as of moderate complexity under CLIA unless the clinical laboratory test or examination is performed under the overall operation and administration of the laboratory director, as described in Section 1209, including, but not limited to, documentation by the laboratory director of the adequacy of the qualifications and competency of the personnel, and the test is performed by any of the following persons:
- (1) A licensed physician and surgeon holding a M.D. or D.O. degree.
 - (2) A licensed podiatrist or a licensed dentist if the results of the tests can be lawfully utilized within his or her practice.
 - (3) A person licensed under this chapter to engage in clinical laboratory practice or to direct a clinical laboratory.
 - (4) A person authorized to perform tests pursuant to a certificate issued under Article 5 (commencing with Section 101150) of Chapter 2 of Part 3 of Division 101 of the Health and Safety Code.
 - (5) A licensed physician assistant if authorized by a supervising physician and surgeon in accordance with Section 3502 or 3535.
 - (6) A person licensed under Chapter 6 (commencing with Section 2700).
 - (7) A perfusionist if authorized by and performed in compliance with Section 2590.
 - (8) A respiratory care practitioner if authorized by and performed in compliance with Chapter 8.3 (commencing with Section 3700).

(9) A person performing nuclear medicine technology if authorized by and performed in compliance with Article 6 (commencing with Section 107150) of Chapter 4 of Part 1 of Division 104 of the Health and Safety Code.

(10) Any person if performing blood gas analysis in compliance with Section 1245.

(11) (A) A person certified or licensed as an "Emergency Medical Technician II" or paramedic pursuant to Division 2.5 (commencing with Section 1797) of the Health and Safety Code while providing prehospital medical care, a person licensed as a psychiatric technician under Chapter 10 (commencing with Section 4500) of Division 2, as a vocational nurse pursuant to Chapter 6.5 (commencing with Section 2840), or as a midwife licensed pursuant to Article 24 (commencing with Section 2505) of Chapter 5, or certified by the department pursuant to Division 5 (commencing with Section 70001) of Title 22 of the California Code of Regulations as a nurse assistant or a home health aide, who provides direct patient care, if the person is performing the test as an adjunct to the provision of direct patient care by the person, is utilizing a point-of-care laboratory testing device at a site for which a laboratory license or registration has been issued, meets the minimum clinical laboratory education, training, and experience requirements set forth in regulations adopted by the department, and has demonstrated to the satisfaction of the laboratory director that he or she is competent in the operation of the point-of-care laboratory testing device for each analyte to be reported.

(B) Prior to being authorized by the laboratory director to perform laboratory tests or examinations, testing personnel identified in subparagraph (A) shall participate in a preceptor program until they are able to perform the clinical laboratory tests or examinations authorized in this section with results that are deemed accurate and skills that are deemed competent by the preceptor. For the purposes of this section, a "preceptor program" means an organized system that meets regulatory requirements in which a preceptor provides and documents personal observation and critical evaluation, including review of accuracy, reliability, and validity, of laboratory testing performed.

(12) Any other person within a physician office laboratory if the test is performed under the supervision of the patient's physician and surgeon or podiatrist who shall be accessible to the laboratory to provide onsite, telephone, or electronic consultation as needed, and shall: (A) ensure that the person is performing test methods as required for accurate and reliable tests; and (B) have personal knowledge of the results of the clinical laboratory testing or examination performed by that person before the test results are reported from the laboratory.

(13) A pharmacist, if ordering drug therapy-related laboratory tests in compliance with paragraph (2) of subdivision (a) of Section 4052.1 or paragraph (2) of subdivision (a) of Section 4052.2.

(c) Notwithstanding subdivision (b) of Section 1206, no person shall perform clinical laboratory tests or examinations classified as of high complexity under CLIA unless the clinical laboratory test or examination is performed under the overall operation and administration of the laboratory director, as described in Section 1209, including, but not limited to, documentation by the laboratory director of the

adequacy of the qualifications and competency of the personnel, and the test is performed by any of the following persons:

- (1) A licensed physician and surgeon holding a M.D. or D.O. degree.
- (2) A licensed podiatrist or a licensed dentist if the results of the tests can be lawfully utilized within his or her practice.
- (3) A person licensed under this chapter to engage in clinical laboratory practice or to direct a clinical laboratory if the test or examination is within a specialty or subspecialty authorized by the person's licensure.
- (4) A person authorized to perform tests pursuant to a certificate issued under Article 5 (commencing with Section 101150) of Chapter 2 of Part 3 of Division 101 of the Health and Safety Code if the test or examination is within a specialty or subspecialty authorized by the person's certification.
- (5) A licensed physician assistant if authorized by a supervising physician and surgeon in accordance with Section 3502 or 3535.
- (6) A perfusionist if authorized by and performed in compliance with Section 2590.
- (7) A respiratory care practitioner if authorized by and performed in compliance with Chapter 8.3 (commencing with Section 3700).
- (8) A person performing nuclear medicine technology if authorized by and performed in compliance with Article 6 (commencing with Section 107150) of Chapter 4 of Part 1 of Division 104 of the Health and Safety Code.
- (9) Any person if performing blood gas analysis in compliance with Section 1245.
- (10) Any other person within a physician office laboratory if the test is performed under the onsite supervision of the patient's physician and surgeon or podiatrist who shall: (A) ensure that the person is performing test methods as required for accurate and reliable tests; and (B) have personal knowledge of the results of clinical laboratory testing or examination performed by that person before the test results are reported from the laboratory.

(d) Clinical laboratory examinations classified as provider-performed microscopy under CLIA may be personally performed using a brightfield or phase/contrast microscope by one of the following practitioners:

- (1) A licensed physician and surgeon using the microscope during the patient's visit on a specimen obtained from his or her own patient or from a patient of a group medical practice of which the physician is a member or employee.
- (2) A nurse midwife holding a certificate as specified by Section 2746.5, a licensed nurse practitioner as specified in Section 2835.5, or a licensed physician assistant acting under the supervision of a physician pursuant to Section 3502 using the microscope during the patient's visit on a specimen obtained from his or her own patient or from the patient of a clinic, group medical practice, or other health care provider of which the certified nurse midwife, licensed nurse practitioner, or licensed physician assistant is an employee.

(3) A licensed dentist using the microscope during the patient's visit on a specimen obtained from his or her own patient or from a patient of a group dental practice of which the dentist is a member or an employee.

Amend BPC section 1209.

(a) As used in this chapter, "laboratory director" means any person who is any of the following:

(1) A duly licensed physician and surgeon.

(2) Only for purposes of a clinical laboratory test or examination classified as waived, is any of the following:

(A) A duly licensed clinical laboratory scientist.

(B) A duly licensed limited clinical laboratory scientist.

(C) A duly licensed naturopathic doctor.

(D) A duly licensed optometrist serving as the director of a laboratory that only performs clinical laboratory tests authorized in paragraph (10) of subdivision (d) of Section 3041.

(E) A pharmacist-in-charge of a pharmacy serving as the director of a laboratory that only performs CLIA waived tests as authorized in Pharmacy Law.

(3) Licensed to direct a clinical laboratory under this chapter.

(b) (1) A person defined in paragraph (1) or (3) of subdivision (a) who is identified as the CLIA laboratory director of a laboratory that performs clinical laboratory tests classified as moderate or high complexity shall also meet the laboratory director qualifications under CLIA for the type and complexity of tests being offered by the laboratory.

(2) As used in this subdivision, "CLIA laboratory director" means the person identified as the laboratory director on the CLIA certificate issued to the laboratory by the federal Centers for Medicare and Medicaid Services (CMS).

(c) The laboratory director, if qualified under CLIA, may perform the duties of the technical consultant, technical supervisor, clinical consultant, general supervisor, and testing personnel, or delegate these responsibilities to persons qualified under CLIA. If the laboratory director reappoints performance of those responsibilities or duties, he or she shall remain responsible for ensuring that all those duties and responsibilities are properly performed.

(d) (1) The laboratory director is responsible for the overall operation and administration of the clinical laboratory, including administering the technical and scientific operation of a clinical laboratory, the selection and supervision of procedures, the reporting of results, and active participation in its operations to the extent necessary to ensure compliance with this act and CLIA. He or she shall be responsible for the proper performance of all laboratory work of all subordinates and shall employ a sufficient number of laboratory personnel with the appropriate education and either experience or training to provide appropriate consultation, properly supervise and accurately perform tests, and report test results in

accordance with the personnel qualifications, duties, and responsibilities described in CLIA and this chapter.

(2) Where a point-of-care laboratory testing device is utilized and provides results for more than one analyte, the testing personnel may perform and report the results of all tests ordered for each analyte for which he or she has been found by the laboratory director to be competent to perform and report.

(e) As part of the overall operation and administration, the laboratory director of a registered laboratory shall document the adequacy of the qualifications (educational background, training, and experience) of the personnel directing and supervising the laboratory and performing the laboratory test procedures and examinations. In determining the adequacy of qualifications, the laboratory director shall comply with any regulations adopted by the department that specify the minimum qualifications for personnel, in addition to any CLIA requirements relative to the education or training of personnel.

(f) As part of the overall operation and administration, the laboratory director of a licensed laboratory shall do all of the following:

(1) Ensure that all personnel, prior to testing biological specimens, have the appropriate education and experience, receive the appropriate training for the type and complexity of the services offered, and have demonstrated that they can perform all testing operations reliably to provide and report accurate results. In determining the adequacy of qualifications, the laboratory director shall comply with any regulations adopted by the department that specify the minimum qualifications for, and the type of procedures that may be performed by, personnel in addition to any CLIA requirements relative to the education or training of personnel. Any regulations adopted pursuant to this section that specify the type of procedure that may be performed by testing personnel shall be based on the skills, knowledge, and tasks required to perform the type of procedure in question.

(2) Ensure that policies and procedures are established for monitoring individuals who conduct preanalytical, analytical, and postanalytical phases of testing to ensure that they are competent and maintain their competency to process biological specimens, perform test procedures, and report test results promptly and proficiently, and, whenever necessary, identify needs for remedial training or continuing education to improve skills.

(3) Specify in writing the responsibilities and duties of each individual engaged in the performance of the preanalytic, analytic, and postanalytic phases of clinical laboratory tests or examinations, including which clinical laboratory tests or examinations the individual is authorized to perform, whether supervision is required for the individual to perform specimen processing, test performance, or results reporting, and whether consultant, supervisor, or director review is required prior to the individual reporting patient test results.

(g) The competency and performance of staff of a licensed laboratory shall be evaluated and documented by the laboratory director, or by a person who qualifies as a technical consultant or a technical supervisor under CLIA depending on the type and complexity of tests being offered by the laboratory.

(1) The procedures for evaluating the competency of the staff shall include, but are not limited to, all of the following:

(A) Direct observations of routine patient test performance, including patient preparation, if applicable, and specimen handling, processing, and testing.

(B) Monitoring the recording and reporting of test results.

(C) Review of intermediate test results or worksheets, quality control records, proficiency testing results, and preventive maintenance records.

(D) Direct observation of performance of instrument maintenance and function checks.

(E) Assessment of test performance through testing previously analyzed specimens, internal blind testing samples, or external proficiency testing samples.

(F) Assessment of problem solving skills.

(2) Evaluation and documentation of staff competency and performance shall occur at least semiannually during the first year an individual tests biological specimens. Thereafter, evaluations shall be performed at least annually unless test methodology or instrumentation changes, in which case, prior to reporting patient test results, the individual's performance shall be reevaluated to include the use of the new test methodology or instrumentation.

(h) The laboratory director of each clinical laboratory of an acute care hospital shall be a physician and surgeon who is a qualified pathologist, except as follows:

(1) If a qualified pathologist is not available, a physician and surgeon or a clinical laboratory bioanalyst qualified as a laboratory director under subdivision (a) may direct the laboratory. However, a qualified pathologist shall be available for consultation at suitable intervals to ensure high-quality service.

(2) If there are two or more clinical laboratories of an acute care hospital, those additional clinical laboratories that are limited to the performance of blood gas analysis, blood electrolyte analysis, or both, may be directed by a physician and surgeon qualified as a laboratory director under subdivision (a), irrespective of whether a pathologist is available.

As used in this subdivision, a qualified pathologist is a physician and surgeon certified or eligible for certification in clinical or anatomical pathology by the American Board of Pathology or the American Osteopathic Board of Pathology.

(i) Subdivision (h) does not apply to any director of a clinical laboratory of an acute care hospital acting in that capacity on or before January 1, 1988.

(j) A laboratory director may serve as the director of up to the maximum number of laboratories stipulated by CLIA, as defined under Section 1202.5.

Attachment 2

BUSINESS AND PROFESSIONS CODE - BPC

DIVISION 2. HEALING ARTS [500 - 4999.129]

(Division 2 enacted by Stats. 1937, Ch. 399.)

CHAPTER 9. Pharmacy [4000 - 4427.8]

(Chapter 9 repealed and added by Stats. 1996, Ch. 890, Sec. 3.)

ARTICLE 7. Pharmacies [4110 - 4126.9]

(Article 7 added by Stats. 1996, Ch. 890, Sec. 3.)

4115.

(a) A pharmacy technician may perform packaging, manipulative, repetitive, or other nondiscretionary tasks only while assisting, and while under the direct supervision and control of, a pharmacist. The pharmacist shall be responsible for the duties performed under his or her supervision by a technician.

(b) This section does not authorize the performance of any tasks specified in subdivision (a) by a pharmacy technician without a pharmacist on duty.

(c) This section does not authorize a pharmacy technician to perform any act requiring the exercise of professional judgment by a pharmacist.

(d) The board shall adopt regulations to specify tasks pursuant to subdivision (a) that a pharmacy technician may perform under the supervision of a pharmacist. Any pharmacy that employs a pharmacy technician shall do so in conformity with the regulations adopted by the board.

(e) A person shall not act as a pharmacy technician without first being licensed by the board as a pharmacy technician.

(f) (1) A pharmacy with only one pharmacist shall have no more than one pharmacy technician performing the tasks specified in subdivision (a). The ratio of pharmacy technicians performing the tasks specified in subdivision (a) to any additional pharmacist shall not exceed 2:1, except that this ratio shall not apply to personnel performing clerical functions pursuant to Section 4116 or 4117. This ratio is applicable to all practice settings, except for an inpatient of a licensed health facility, a patient of a licensed home health agency, as specified in paragraph (2), an inmate of a correctional facility of the Department of Corrections and Rehabilitation, and for a person receiving treatment in a facility

operated by the State Department of State Hospitals, the State Department of Developmental Services, or the Department of Veterans Affairs.

(2) The board may adopt regulations establishing the ratio of pharmacy technicians performing the tasks specified in subdivision (a) to pharmacists applicable to the filling of prescriptions of an inpatient of a licensed health facility and for a patient of a licensed home health agency. Any ratio established by the board pursuant to this subdivision shall allow, at a minimum, at least one pharmacy technician for a single pharmacist in a pharmacy and two pharmacy technicians for each additional pharmacist, except that this ratio shall not apply to personnel performing clerical functions pursuant to Section 4116 or 4117.

(3) A pharmacist scheduled to supervise a second pharmacy technician may refuse to supervise a second pharmacy technician if the pharmacist determines, in the exercise of his or her professional judgment, that permitting the second pharmacy technician to be on duty would interfere with the effective performance of the pharmacist's responsibilities under this chapter. A pharmacist assigned to supervise a second pharmacy technician shall notify the pharmacist in charge in writing of his or her determination, specifying the circumstances of concern with respect to the pharmacy or the pharmacy technician that have led to the determination, within a reasonable period, but not to exceed 24 hours, after the posting of the relevant schedule. An entity employing a pharmacist shall not discharge, discipline, or otherwise discriminate against any pharmacist in the terms and conditions of employment for exercising or attempting to exercise in good faith the right established pursuant to this paragraph.

(g) Notwithstanding subdivisions (a) and (b), the board shall by regulation establish conditions to permit the temporary absence of a pharmacist for breaks and lunch periods pursuant to Section 512 of the Labor Code and the orders of the Industrial Welfare Commission without closing the pharmacy. During these temporary absences, a pharmacy technician may, at the discretion of the pharmacist, remain in the pharmacy but may only perform nondiscretionary tasks. The pharmacist shall be responsible for a pharmacy technician and shall review any task performed by a pharmacy technician during the pharmacist's temporary absence. This subdivision shall not be construed to authorize a pharmacist to supervise pharmacy technicians in greater ratios than those described in subdivision (f).

(h) The pharmacist on duty shall be directly responsible for the conduct of a pharmacy technician supervised by that pharmacist.

(i) In a health care facility licensed under subdivision (a) of Section 1250 of the Health and Safety Code, a pharmacy technician's duties may include any of the following:

(1) Packaging emergency supplies for use in the health care facility and the hospital's emergency medical system or as authorized under Section 4119.

(2) Sealing emergency containers for use in the health care facility.

(3) Performing monthly checks of the drug supplies stored throughout the health care facility. Irregularities shall be reported within 24 hours to the pharmacist in charge and the director or chief executive officer of the health care facility in accordance with the health care facility's policies and procedures.

(Amended by Stats. 2015, Ch. 303, Sec. 5. (AB 731) Effective January 1, 2016.)

§ 1793.2. Duties of a Pharmacy Technician.

16 CA ADC § 1793.2 BARCLAYS OFFICIAL CALIFORNIA CODE OF REGULATIONS

Barclays Official California Code of Regulations

Title 16. Professional and Vocational Regulations

Division 17. California State Board of Pharmacy

Article 11. Ancillary Personnel (Refs & Annos)

16 CCR § 1793.2

§ 1793.2. Duties of a Pharmacy Technician.

“Nondiscretionary tasks” as used in Business and Professions Code section 4115, include:

- (a) removing the drug or drugs from stock;
- (b) counting, pouring, or mixing pharmaceuticals;
- (c) placing the product into a container;
- (d) affixing the label or labels to the container;
- (e) packaging and repackaging.

Note: Authority cited: Sections 4005, 4007, 4038, 4115 and 4202, Business and Professions Code. Reference: Sections 4005, 4007, 4038, 4115 and 4202, Business and Professions Code.

HISTORY

1. New section filed 8-12-92; operative 9-11-92 (Register 92, No. 33).
2. Change without regulatory effect amending first paragraph and Note filed 9-11-2002 pursuant to section 100, title 1, California Code of Regulations (Register 2002, No. 37).
3. Repealer of first paragraph filed 9-22-2004; operative 10-22-2004 (Register 2004, No. 39).

This database is current through 1/8/21 Register 2021, No. 2

16 CCR § 1793.2, 16 CA ADC § 1793.2

Proposal to Amend Business and Professions Code section 4115.

(a) A pharmacy technician may perform packaging, manipulative, repetitive, or other nondiscretionary tasks only while assisting, and while under the direct supervision and control of, a pharmacist. The pharmacist shall be responsible for the duties performed under his or her supervision by a technician.

(b) This section does not authorize the performance of any tasks specified in subdivision (a) by a pharmacy technician without a pharmacist on duty.

(c) This section does not authorize a pharmacy technician to perform any act requiring the exercise of professional judgment by a pharmacist.

(d) The board shall adopt regulations to specify tasks pursuant to subdivision (a) that a pharmacy technician may perform under the supervision of a pharmacist. Any pharmacy that employs a pharmacy technician shall do so in conformity with the regulations adopted by the board.

(e) A person shall not act as a pharmacy technician without first being licensed by the board as a pharmacy technician.

(f) (1) A pharmacy with only one pharmacist shall have no more than one pharmacy technician performing the tasks specified in subdivision (a). The ratio of pharmacy technicians performing the tasks specified in subdivision (a) to any additional pharmacist shall not exceed 2:1, except that this ratio shall not apply to personnel performing clerical functions pursuant to Section 4116 or 4117. This ratio is applicable to all practice settings, except for an inpatient of a licensed health facility, a patient of a licensed home health agency, as specified in paragraph (2), an inmate of a correctional facility of the Department of Corrections and Rehabilitation, and for a person receiving treatment in a facility operated by the State Department of State Hospitals, the State Department of Developmental Services, or the Department of Veterans Affairs.

(2) The board may adopt regulations establishing the ratio of pharmacy technicians performing the tasks specified in subdivision (a) to pharmacists applicable to the filling of prescriptions of an inpatient of a licensed health facility and for a patient of a licensed home health agency. Any ratio established by the board pursuant to this subdivision shall allow, at a minimum, at least one pharmacy technician for a single pharmacist in a pharmacy and two pharmacy technicians for each additional pharmacist, except that this ratio shall not apply to personnel performing clerical functions pursuant to Section 4116 or 4117.

(3) A pharmacist scheduled to supervise a second pharmacy technician may refuse to supervise a second pharmacy technician if the pharmacist determines, in the exercise of his or her professional judgment, that permitting the second pharmacy technician to be on duty would interfere with the effective performance of the pharmacist's responsibilities under this chapter. A pharmacist assigned to supervise a second pharmacy technician shall notify the pharmacist in charge in writing of his or her determination, specifying the circumstances of concern with respect to the pharmacy or the pharmacy technician that have led to the determination, within a reasonable period, but not to exceed 24 hours, after the posting of the relevant schedule. An entity employing a pharmacist shall not discharge, discipline, or otherwise discriminate

against any pharmacist in the terms and conditions of employment for exercising or attempting to exercise in good faith the right established pursuant to this paragraph.

(g) Notwithstanding subdivisions (a) and (b), the board shall by regulation establish conditions to permit the temporary absence of a pharmacist for breaks and lunch periods pursuant to Section 512 of the Labor Code and the orders of the Industrial Welfare Commission without closing the pharmacy. During these temporary absences, a pharmacy technician may, at the discretion of the pharmacist, remain in the pharmacy but may only perform nondiscretionary tasks. The pharmacist shall be responsible for a pharmacy technician and shall review any task performed by a pharmacy technician during the pharmacist's temporary absence. This subdivision shall not be construed to authorize a pharmacist to supervise pharmacy technicians in greater ratios than those described in subdivision (f).

(h) The pharmacist on duty shall be directly responsible for the conduct of a pharmacy technician supervised by that pharmacist.

(i) In a health care facility licensed under subdivision (a) of Section 1250 of the Health and Safety Code, a pharmacy technician's duties may include any of the following:

(1) Packaging emergency supplies for use in the health care facility and the hospital's emergency medical system or as authorized under Section 4119.

(2) Sealing emergency containers for use in the health care facility.

(3) Performing monthly checks of the drug supplies stored throughout the health care facility. Irregularities shall be reported within 24 hours to the pharmacist in charge and the director or chief executive officer of the health care facility in accordance with the health care facility's policies and procedures.

(j) A pharmacy technician may administer a COVID-19 or influenza vaccine, if deemed appropriate and delegated by the supervising pharmacist, if the following conditions are met:

1. The pharmacy technician holds a current certification in cardiopulmonary resuscitation (CPR).
2. The pharmacy technician has completed a training consisting of a minimum of 6 hours of training, which includes live training, assessment and evaluation of injection technique assessment, and completing the online assessment and evaluation from an ACPE accredited provider.
3. The pharmacy technician completes 1 hours of immunization related continuing education once every two years.
4. If deemed appropriate by the supervising pharmacist, a pharmacy technician may also administer epinephrine.
5. The pharmacy maintains a record of the identification of the pharmacy technician administering the vaccine and the identification of the supervising pharmacist.

Attachment 3

Recommended Child and Adolescent Immunization Schedule for ages 18 years or younger

UNITED STATES
2020

Vaccines in the Child and Adolescent Immunization Schedule*

Vaccines	Abbreviations	Trade names
Diphtheria, tetanus, and acellular pertussis vaccine	DTaP	Daptacel® Infanrix®
Diphtheria, tetanus vaccine	DT	No trade name
<i>Haemophilus influenzae</i> type b vaccine	Hib (PRP-T) Hib (PRP-OMP)	ActHIB® Hiberix® PedvaxHIB®
Hepatitis A vaccine	HepA	Havrix® Vaqta®
Hepatitis B vaccine	HepB	Engerix-B® Recombivax HB®
Human papillomavirus vaccine	HPV	Gardasil 9®
Influenza vaccine (inactivated)	IIV	Multiple
Influenza vaccine (live, attenuated)	LAIV	FluMist® Quadrivalent
Measles, mumps, and rubella vaccine	MMR	M-M-R® II
Meningococcal serogroups A, C, W, Y vaccine	MenACWY-D	Menactra®
	MenACWY-CRM	Menveo®
Meningococcal serogroup B vaccine	MenB-4C MenB-FHbp	Bexsero® Trumenba®
Pneumococcal 13-valent conjugate vaccine	PCV13	Prevnar 13®
Pneumococcal 23-valent polysaccharide vaccine	PPSV23	Pneumovax® 23
Poliovirus vaccine (inactivated)	IPV	IPOL®
Rotavirus vaccine	RV1 RV5	Rotarix® RotaTeq®
Tetanus, diphtheria, and acellular pertussis vaccine	Tdap	Adacel® Boostrix®
Tetanus and diphtheria vaccine	Td	Tenivac® Tdvax™
Varicella vaccine	VAR	Varivax®
Combination vaccines (use combination vaccines instead of separate injections when appropriate)		
DTaP, hepatitis B, and inactivated poliovirus vaccine	DTaP-HepB-IPV	Pediarix®
DTaP, inactivated poliovirus, and <i>Haemophilus influenzae</i> type b vaccine	DTaP-IPV/Hib	Pentacel®
DTaP and inactivated poliovirus vaccine	DTaP-IPV	Kinrix® Quadracel®
Measles, mumps, rubella, and varicella vaccine	MMRV	ProQuad®

*Administer recommended vaccines if immunization history is incomplete or unknown. Do not restart or add doses to vaccine series for extended intervals between doses. When a vaccine is not administered at the recommended age, administer at a subsequent visit. The use of trade names is for identification purposes only and does not imply endorsement by the ACIP or CDC.

How to use the child/adolescent immunization schedule

- 1** Determine recommended vaccine by age (**Table 1**)
- 2** Determine recommended interval for catch-up vaccination (**Table 2**)
- 3** Assess need for additional recommended vaccines by medical condition and other indications (**Table 3**)
- 4** Review vaccine types, frequencies, intervals, and considerations for special situations (**Notes**)

Recommended by the Advisory Committee on Immunization Practices (www.cdc.gov/vaccines/acip) and approved by the Centers for Disease Control and Prevention (www.cdc.gov), American Academy of Pediatrics (www.aap.org), American Academy of Family Physicians (www.aafp.org), American College of Obstetricians and Gynecologists (www.acog.org), and American College of Nurse-Midwives (www.midwife.org).

Report

- Suspected cases of reportable vaccine-preventable diseases or outbreaks to your state or local health department
- Clinically significant adverse events to the Vaccine Adverse Event Reporting System (VAERS) at www.vaers.hhs.gov or 800-822-7967



Download the CDC Vaccine Schedules App for providers at www.cdc.gov/vaccines/schedules/hcp/schedule-app.html.

Helpful information

- Complete ACIP recommendations: www.cdc.gov/vaccines/hcp/acip-recs/index.html
- General Best Practice Guidelines for Immunization: www.cdc.gov/vaccines/hcp/acip-recs/general-recs/index.html
- Outbreak information (including case identification and outbreak response), see Manual for the Surveillance of Vaccine-Preventable Diseases: www.cdc.gov/vaccines/pubs/surv-manual



U.S. Department of Health and Human Services
Centers for Disease Control and Prevention

Table 1

Recommended Child and Adolescent Immunization Schedule for ages 18 years or younger, United States, 2020

These recommendations must be read with the notes that follow. For those who fall behind or start late, provide catch-up vaccination at the earliest opportunity as indicated by the green bars. To determine minimum intervals between doses, see the catch-up schedule (Table 2). School entry and adolescent vaccine age groups are shaded in gray.

Vaccine	Birth	1 mo	2 mos	4 mos	6 mos	9 mos	12 mos	15 mos	18 mos	19–23 mos	2–3 yrs	4–6 yrs	7–10 yrs	11–12 yrs	13–15 yrs	16 yrs	17–18 yrs	
Hepatitis B (HepB)	1 st dose	2 nd dose			←----- 3 rd dose -----→													
Rotavirus (RV): RV1 (2-dose series), RV5 (3-dose series)			1 st dose	2 nd dose	See Notes													
Diphtheria, tetanus, acellular pertussis (DTaP <7 yrs)			1 st dose	2 nd dose	3 rd dose			←----- 4 th dose -----→				5 th dose						
Haemophilus influenzae type b (Hib)			1 st dose	2 nd dose	See Notes		← 3 rd or 4 th dose, See Notes →											
Pneumococcal conjugate (PCV13)			1 st dose	2 nd dose	3 rd dose		←----- 4 th dose -----→											
Inactivated poliovirus (IPV <18 yrs)			1 st dose	2 nd dose	←----- 3 rd dose -----→						4 th dose							
Influenza (IIV)					Annual vaccination 1 or 2 doses								Annual vaccination 1 dose only					
Influenza (LAIV)												Annual vaccination 1 or 2 doses		Annual vaccination 1 dose only				
Measles, mumps, rubella (MMR)					See Notes	←----- 1 st dose -----→						2 nd dose						
Varicella (VAR)						←----- 1 st dose -----→						2 nd dose						
Hepatitis A (HepA)					See Notes	2-dose series, See Notes												
Tetanus, diphtheria, acellular pertussis (Tdap ≥7 yrs)																		Tdap
Human papillomavirus (HPV)														*	See Notes			
Meningococcal (MenACWY-D ≥9 mos, MenACWY-CRM ≥2 mos)			See Notes											1 st dose		2 nd dose		
Meningococcal B															See Notes			
Pneumococcal polysaccharide (PPSV23)												See Notes						

Range of recommended ages for all children
 Range of recommended ages for catch-up immunization
 Range of recommended ages for certain high-risk groups
 Recommended based on shared clinical decision-making or *can be used in this age group
 No recommendation/ not applicable

Table 2

Recommended Catch-up Immunization Schedule for Children and Adolescents Who Start Late or Who are More than 1 month Behind, United States, 2020

The table below provides catch-up schedules and minimum intervals between doses for children whose vaccinations have been delayed. A vaccine series does not need to be restarted, regardless of the time that has elapsed between doses. Use the section appropriate for the child's age. **Always use this table in conjunction with Table 1 and the notes that follow.**

Children age 4 months through 6 years					
Vaccine	Minimum Age for Dose 1	Minimum Interval Between Doses			
		Dose 1 to Dose 2	Dose 2 to Dose 3	Dose 3 to Dose 4	Dose 4 to Dose 5
Hepatitis B	Birth	4 weeks	8 weeks and at least 16 weeks after first dose. Minimum age for the final dose is 24 weeks.		
Rotavirus	6 weeks Maximum age for first dose is 14 weeks, 6 days	4 weeks	4 weeks Maximum age for final dose is 8 months, 0 days.		
Diphtheria, tetanus, and acellular pertussis	6 weeks	4 weeks	4 weeks	6 months	6 months
<i>Haemophilus influenzae</i> type b	6 weeks	No further doses needed if first dose was administered at age 15 months or older. 4 weeks if first dose was administered before the 1 st birthday. 8 weeks (as final dose) if first dose was administered at age 12 through 14 months.	No further doses needed if previous dose was administered at age 15 months or older. 4 weeks if current age is younger than 12 months and first dose was administered at younger than age 7 months and at least 1 previous dose was PRP-T (ActHib, Pentacel, Hiberix) or unknown. 8 weeks and age 12 through 59 months (as final dose) if current age is younger than 12 months and first dose was administered at age 7 through 11 months; OR if current age is 12 through 59 months and first dose was administered before the 1 st birthday and second dose administered at younger than 15 months; OR if both doses were PRP-OMP (PedvaxHIB, Comvax) and were administered before the 1 st birthday.	8 weeks (as final dose) This dose only necessary for children age 12 through 59 months who received 3 doses before the 1 st birthday.	
Pneumococcal conjugate	6 weeks	No further doses needed for healthy children if first dose was administered at age 24 months or older. 4 weeks if first dose was administered before the 1 st birthday. 8 weeks (as final dose for healthy children) if first dose was administered at the 1 st birthday or after.	No further doses needed for healthy children if previous dose administered at age 24 months or older. 4 weeks if current age is younger than 12 months and previous dose was administered at <7 months old. 8 weeks (as final dose for healthy children) if previous dose was administered between 7–11 months (wait until at least 12 months old); OR if current age is 12 months or older and at least 1 dose was given before age 12 months.	8 weeks (as final dose) This dose only necessary for children age 12 through 59 months who received 3 doses before age 12 months or for children at high risk who received 3 doses at any age.	
Inactivated poliovirus	6 weeks	4 weeks	4 weeks if current age is < 4 years. 6 months (as final dose) if current age is 4 years or older.	6 months (minimum age 4 years for final dose).	
Measles, mumps, rubella	12 months	4 weeks			
Varicella	12 months	3 months			
Hepatitis A	12 months	6 months			
Meningococcal ACWY	2 months MenACWY-CRM 9 months MenACWY-D	8 weeks	See Notes	See Notes	
Children and adolescents age 7 through 18 years					
Meningococcal ACWY	Not applicable (N/A)	8 weeks			
Tetanus, diphtheria; tetanus, diphtheria, and acellular pertussis	7 years	4 weeks	4 weeks if first dose of DTaP/DT was administered before the 1 st birthday. 6 months (as final dose) if first dose of DTaP/DT or Tdap/Td was administered at or after the 1 st birthday.	6 months if first dose of DTaP/DT was administered before the 1 st birthday.	
Human papillomavirus	9 years	Routine dosing intervals are recommended.			
Hepatitis A	N/A	6 months			
Hepatitis B	N/A	4 weeks	8 weeks and at least 16 weeks after first dose.		
Inactivated poliovirus	N/A	4 weeks	6 months A fourth dose is not necessary if the third dose was administered at age 4 years or older and at least 6 months after the previous dose.	A fourth dose of IPV is indicated if all previous doses were administered at <4 years or if the third dose was administered <6 months after the second dose.	
Measles, mumps, rubella	N/A	4 weeks			
Varicella	N/A	3 months if younger than age 13 years. 4 weeks if age 13 years or older.			

Table 3

Recommended Child and Adolescent Immunization Schedule by Medical Indication, United States, 2020

Always use this table in conjunction with Table 1 and the notes that follow.

VACCINE	INDICATION									
	Pregnancy	Immunocompromised status (excluding HIV infection)	HIV infection CD4+ count ¹		Kidney failure, end-stage renal disease, or on hemodialysis	Heart disease or chronic lung disease	CSF leaks or cochlear implants	Asplenia or persistent complement deficiencies	Chronic liver disease	Diabetes
			<15% and total CD4 cell count of <200/mm ³	≥15% and total CD4 cell count of ≥200/mm ³						
Hepatitis B										
Rotavirus		SCID ²								
Diphtheria, tetanus, & acellular pertussis (DTaP)										
<i>Haemophilus influenzae</i> type b										
Pneumococcal conjugate										
Inactivated poliovirus										
Influenza (IIV)										
Influenza (LAIV)						Asthma, wheezing: 2–4yrs ³				
Measles, mumps, rubella										
Varicella										
Hepatitis A										
Tetanus, diphtheria, & acellular pertussis (Tdap)										
Human papillomavirus										
Meningococcal ACWY										
Meningococcal B										
Pneumococcal polysaccharide										

Vaccination according to the routine schedule recommended
 Recommended for persons with an additional risk factor for which the vaccine would be indicated
 Vaccination is recommended, and additional doses may be necessary based on medical condition. See Notes.
 Not recommended/contraindicated—vaccine should not be administered
 Precaution—vaccine might be indicated if benefit of protection outweighs risk of adverse reaction
 Delay vaccination until after pregnancy if vaccine indicated
 No recommendation/not applicable

¹ For additional information regarding HIV laboratory parameters and use of live vaccines, see the General Best Practice Guidelines for Immunization, “Altered Immunocompetence,” at www.cdc.gov/vaccines/hcp/acip-recs/general-recs/immunocompetence.html and Table 4-1 (footnote D) at www.cdc.gov/vaccines/hcp/acip-recs/general-recs/contraindications.html.

² Severe Combined Immunodeficiency

³ LAIV contraindicated for children 2–4 years of age with asthma or wheezing during the preceding 12 months.

For vaccine recommendations for persons 19 years of age or older, see the Recommended Adult Immunization Schedule.

Additional information

- Consult relevant ACIP statements for detailed recommendations at www.cdc.gov/vaccines/hcp/acip-recs/index.html.
- For information on contraindications and precautions for the use of a vaccine, consult the General Best Practice Guidelines for Immunization at www.cdc.gov/vaccines/hcp/acip-recs/general-recs/contraindications.html and relevant ACIP statements at www.cdc.gov/vaccines/hcp/acip-recs/index.html.
- For calculating intervals between doses, 4 weeks = 28 days. Intervals of ≥ 4 months are determined by calendar months.
- Within a number range (e.g., 12–18), a dash (–) should be read as “through.”
- Vaccine doses administered ≤ 4 days before the minimum age or interval are considered valid. Doses of any vaccine administered ≥ 5 days earlier than the minimum age or minimum interval should not be counted as valid and should be repeated as age-appropriate. The repeat dose should be spaced after the invalid dose by the recommended minimum interval. For further details, see Table 3-1, Recommended and minimum ages and intervals between vaccine doses, in General Best Practice Guidelines for Immunization at www.cdc.gov/vaccines/hcp/acip-recs/general-recs/timing.html.
- Information on travel vaccine requirements and recommendations is available at www.cdc.gov/travel/.
- For vaccination of persons with immunodeficiencies, see Table 8-1, Vaccination of persons with primary and secondary immunodeficiencies, in General Best Practice Guidelines for Immunization at www.cdc.gov/vaccines/hcp/acip-recs/general-recs/immunocompetence.html, and Immunization in Special Clinical Circumstances (In: Kimberlin DW, Brady MT, Jackson MA, Long SS, eds. *Red Book: 2018 Report of the Committee on Infectious Diseases*. 31st ed. Itasca, IL: American Academy of Pediatrics; 2018:67–111).
- For information regarding vaccination in the setting of a vaccine-preventable disease outbreak, contact your state or local health department.
- The National Vaccine Injury Compensation Program (VICP) is a no-fault alternative to the traditional legal system for resolving vaccine injury claims. All routine child and adolescent vaccines are covered by VICP except for pneumococcal polysaccharide vaccine (PPSV23). For more information, see www.hrsa.gov/vaccinecompensation/index.html.

Diphtheria, tetanus, and pertussis (DTaP) vaccination (minimum age: 6 weeks [4 years for Kinrix or Quadracel])

Routine vaccination

- 5-dose series at 2, 4, 6, 15–18 months, 4–6 years
 - **Prospectively:** Dose 4 may be administered as early as age 12 months if at least 6 months have elapsed since dose 3.
 - **Retrospectively:** A 4th dose that was inadvertently administered as early as 12 months may be counted if at least 4 months have elapsed since dose 3.

Catch-up vaccination

- Dose 5 is not necessary if dose 4 was administered at age 4 years or older and at least 6 months after dose 3.
- For other catch-up guidance, see Table 2.

Haemophilus influenzae type b vaccination (minimum age: 6 weeks)

Routine vaccination

- **ActHIB, Hiberix, or Pentacel:** 4-dose series at 2, 4, 6, 12–15 months
- **PedvaxHIB:** 3-dose series at 2, 4, 12–15 months

Catch-up vaccination

- **Dose 1 at 7–11 months:** Administer dose 2 at least 4 weeks later and dose 3 (final dose) at 12–15 months or 8 weeks after dose 2 (whichever is later).
- **Dose 1 at 12–14 months:** Administer dose 2 (final dose) at least 8 weeks after dose 1.
- **Dose 1 before 12 months and dose 2 before 15 months:** Administer dose 3 (final dose) 8 weeks after dose 2.
- **2 doses of PedvaxHIB before 12 months:** Administer dose 3 (final dose) at 12–59 months and at least 8 weeks after dose 2.
- **Unvaccinated at 15–59 months:** 1 dose
- **Previously unvaccinated children age 60 months or older** who are not considered high risk do not require catch-up vaccination.
- For other catch-up guidance, see Table 2.

Special situations

• Chemotherapy or radiation treatment:

12–59 months

- Unvaccinated or only 1 dose before age 12 months: 2 doses, 8 weeks apart
- 2 or more doses before age 12 months: 1 dose at least 8 weeks after previous dose

Doses administered within 14 days of starting therapy or during therapy should be repeated at least 3 months after therapy completion.

• Hematopoietic stem cell transplant (HSCT):

- 3-dose series 4 weeks apart starting 6 to 12 months after successful transplant, regardless of Hib vaccination history

• Anatomic or functional asplenia (including sickle cell disease):

12–59 months

- Unvaccinated or only 1 dose before age 12 months: 2 doses, 8 weeks apart
- 2 or more doses before age 12 months: 1 dose at least 8 weeks after previous dose

Unvaccinated* persons age 5 years or older

- 1 dose

• Elective splenectomy:

Unvaccinated* persons age 15 months or older

- 1 dose (preferably at least 14 days before procedure)

• HIV infection:

12–59 months

- Unvaccinated or only 1 dose before age 12 months: 2 doses, 8 weeks apart
- 2 or more doses before age 12 months: 1 dose at least 8 weeks after previous dose

Unvaccinated* persons age 5–18 years

- 1 dose

• Immunoglobulin deficiency, early component complement deficiency:

12–59 months

- Unvaccinated or only 1 dose before age 12 months: 2 doses, 8 weeks apart
- 2 or more doses before age 12 months: 1 dose at least 8 weeks after previous dose

*Unvaccinated = Less than routine series (through 14 months) OR no doses (15 months or older)

Notes

Recommended Child and Adolescent Immunization Schedule for ages 18 years or younger, United States, 2020

Hepatitis A vaccination

(minimum age: 12 months for routine vaccination)

Routine vaccination

- 2-dose series (minimum interval: 6 months) beginning at age 12 months

Catch-up vaccination

- Unvaccinated persons through 18 years should complete a 2-dose series (minimum interval: 6 months).
- Persons who previously received 1 dose at age 12 months or older should receive dose 2 at least 6 months after dose 1.
- Adolescents 18 years and older may receive the combined HepA and HepB vaccine, **Twinrix**[®], as a 3-dose series (0, 1, and 6 months) or 4-dose series (0, 7, and 21–30 days, followed by a dose at 12 months).

International travel

- Persons traveling to or working in countries with high or intermediate endemic hepatitis A (www.cdc.gov/travel/):
 - **Infants age 6–11 months:** 1 dose before departure; revaccinate with 2 doses, separated by at least 6 months, between 12 and 23 months of age
 - **Unvaccinated age 12 months and older:** Administer dose 1 as soon as travel is considered.

Hepatitis B vaccination

(minimum age: birth)

Birth dose (monovalent HepB vaccine only)

- **Mother is HBsAg-negative:** 1 dose within 24 hours of birth for **all** medically stable infants $\geq 2,000$ grams. Infants $< 2,000$ grams: Administer 1 dose at chronological age 1 month or hospital discharge.
- **Mother is HBsAg-positive:**
 - Administer **HepB vaccine** and **hepatitis B immune globulin (HBIG)** (in separate limbs) within 12 hours of birth, regardless of birth weight. For infants $< 2,000$ grams, administer 3 additional doses of vaccine (total of 4 doses) beginning at age 1 month.
 - Test for HBsAg and anti-HBs at age 9–12 months. If HepB series is delayed, test 1–2 months after final dose.
- **Mother's HBsAg status is unknown:**
 - Administer **HepB vaccine** within 12 hours of birth, regardless of birth weight.
 - For infants $< 2,000$ grams, administer **HBIG** in addition to HepB vaccine (in separate limbs) within 12 hours of birth. Administer 3 additional doses of vaccine (total of 4 doses) beginning at age 1 month.
 - Determine mother's HBsAg status as soon as possible. If mother is HBsAg-positive, administer **HBIG** to infants $\geq 2,000$ grams as soon as possible, but no later than 7 days of age.

Routine series

- 3-dose series at 0, 1–2, 6–18 months (use monovalent HepB vaccine for doses administered before age 6 weeks)

- Infants who did not receive a birth dose should begin the series as soon as feasible (see Table 2).
- Administration of **4 doses** is permitted when a combination vaccine containing HepB is used after the birth dose.
- **Minimum age** for the final (3rd or 4th) dose: 24 weeks
- **Minimum intervals:** dose 1 to dose 2: 4 weeks / dose 2 to dose 3: 8 weeks / dose 1 to dose 3: 16 weeks (when 4 doses are administered, substitute “dose 4” for “dose 3” in these calculations)

Catch-up vaccination

- Unvaccinated persons should complete a 3-dose series at 0, 1–2, 6 months.
- Adolescents age 11–15 years may use an alternative 2-dose schedule with at least 4 months between doses (adult formulation **Recombivax HB** only).
- Adolescents 18 years and older may receive a 2-dose series of HepB (**Heplisav-B**[®]) at least 4 weeks apart.
- Adolescents 18 years and older may receive the combined HepA and HepB vaccine, **Twinrix**, as a 3-dose series (0, 1, and 6 months) or 4-dose series (0, 7, and 21–30 days, followed by a dose at 12 months).
- For other catch-up guidance, see Table 2.

Special situations

- Revaccination is not generally recommended for persons with a normal immune status who were vaccinated as infants, children, adolescents, or adults.
- **Revaccination** may be recommended for certain populations, including:
 - **Infants born to HBsAg-positive mothers**
 - **Hemodialysis patients**
 - **Other immunocompromised persons**
- For detailed revaccination recommendations, see www.cdc.gov/vaccines/hcp/acip-recs/vacc-specific/hepb.html.

Human papillomavirus vaccination

(minimum age: 9 years)

Routine and catch-up vaccination

- HPV vaccination routinely recommended at **age 11–12 years (can start at age 9 years)** and catch-up HPV vaccination recommended for all persons through age 18 years if not adequately vaccinated
- 2- or 3-dose series depending on age at initial vaccination:
 - **Age 9 through 14 years at initial vaccination:** 2-dose series at 0, 6–12 months (minimum interval: 5 months; repeat dose if administered too soon)
 - **Age 15 years or older at initial vaccination:** 3-dose series at 0, 1–2 months, 6 months (minimum intervals: dose 1 to dose 2: 4 weeks / dose 2 to dose 3: 12 weeks / dose 1 to dose 3: 5 months; repeat dose if administered too soon)
- If completed valid vaccination series with any HPV vaccine, no additional doses needed

Special situations

- **Immunocompromising conditions, including HIV infection:** 3-dose series as above
- **History of sexual abuse or assault:** Start at age 9 years.
- **Pregnancy:** HPV vaccination not recommended until after pregnancy; no intervention needed if vaccinated while pregnant; pregnancy testing not needed before vaccination

Influenza vaccination

(minimum age: 6 months [IIV], 2 years [LAIV], 18 years [recombinant influenza vaccine, RIV])

Routine vaccination

- Use any influenza vaccine appropriate for age and health status annually:
 - 2 doses, separated by at least 4 weeks, for **children age 6 months–8 years** who have received fewer than 2 influenza vaccine doses before July 1, 2019, or whose influenza vaccination history is unknown (administer dose 2 even if the child turns 9 between receipt of dose 1 and dose 2)
 - 1 dose for **children age 6 months–8 years** who have received at least 2 influenza vaccine doses before July 1, 2019
 - 1 dose for **all persons age 9 years and older**
- For the 2020–21 season, see the 2020–21 ACIP influenza vaccine recommendations.

Special situations

- **Egg allergy, hives only:** Any influenza vaccine appropriate for age and health status annually
- **Egg allergy with symptoms other than hives** (e.g., angioedema, respiratory distress, need for emergency medical services or epinephrine): Any influenza vaccine appropriate for age and health status annually in medical setting under supervision of health care provider who can recognize and manage severe allergic reactions
- **LAIV should not be used** in persons with the following conditions or situations:
 - History of severe allergic reaction to a previous dose of any influenza vaccine or to any vaccine component (excluding egg, see details above)
 - Receiving aspirin or salicylate-containing medications
 - Age 2–4 years with history of asthma or wheezing
 - Immunocompromised due to any cause (including medications and HIV infection)
 - Anatomic or functional asplenia
 - Cochlear implant
 - Cerebrospinal fluid-opharyngeal communication
 - Close contacts or caregivers of severely immunosuppressed persons who require a protected environment
 - Pregnancy
 - Received influenza antiviral medications within the previous 48 hours

Measles, mumps, and rubella vaccination (minimum age: 12 months for routine vaccination)

Routine vaccination

- 2-dose series at 12–15 months, 4–6 years
- Dose 2 may be administered as early as 4 weeks after dose 1.

Catch-up vaccination

- Unvaccinated children and adolescents: 2-dose series at least 4 weeks apart
- The maximum age for use of MMRV is 12 years.

Special situations

International travel

- **Infants age 6–11 months:** 1 dose before departure; revaccinate with 2-dose series with dose 1 at 12–15 months (12 months for children in high-risk areas) and dose 2 as early as 4 weeks later.
- **Unvaccinated children age 12 months and older:** 2-dose series at least 4 weeks apart before departure

Meningococcal serogroup A,C,W,Y vaccination (minimum age: 2 months [MenACWY-CRM, Menveo], 9 months [MenACWY-D, Menactra])

Routine vaccination

- 2-dose series at 11–12 years, 16 years

Catch-up vaccination

- Age 13–15 years: 1 dose now and booster at age 16–18 years (minimum interval: 8 weeks)
- Age 16–18 years: 1 dose

Special situations

Anatomic or functional asplenia (including sickle cell disease), HIV infection, persistent complement component deficiency, complement inhibitor (e.g., eculizumab, ravulizumab) use:

- **Menveo**
 - Dose 1 at age 8 weeks: 4-dose series at 2, 4, 6, 12 months
 - Dose 1 at age 7–23 months: 2-dose series (dose 2 at least 12 weeks after dose 1 and after age 12 months)
 - Dose 1 at age 24 months or older: 2-dose series at least 8 weeks apart
- **Menactra**
 - **Persistent complement component deficiency or complement inhibitor use:**
 - Age 9–23 months: 2-dose series at least 12 weeks apart
 - Age 24 months or older: 2-dose series at least 8 weeks apart
 - **Anatomic or functional asplenia, sickle cell disease, or HIV infection:**
 - Age 9–23 months: Not recommended
 - Age 24 months or older: 2-dose series at least 8 weeks apart
 - **Menactra** must be administered at least 4 weeks after completion of PCV13 series.

Travel in countries with hyperendemic or epidemic meningococcal disease, including countries in the African meningitis belt or during the Hajj (www.cdc.gov/travel/):

- Children less than age 24 months:
 - **Menveo (age 2–23 months):**
 - Dose 1 at 8 weeks: 4-dose series at 2, 4, 6, 12 months
 - Dose 1 at 7–23 months: 2-dose series (dose 2 at least 12 weeks after dose 1 and after age 12 months)
 - **Menactra (age 9–23 months):**
 - 2-dose series (dose 2 at least 12 weeks after dose 1; dose 2 may be administered as early as 8 weeks after dose 1 in travelers)
- Children age 2 years or older: 1 dose **Menveo** or **Menactra**

First-year college students who live in residential housing (if not previously vaccinated at age 16 years or older) or military recruits:

- 1 dose **Menveo** or **Menactra**

Adolescent vaccination of children who received MenACWY prior to age 10 years:

- **Children for whom boosters are recommended** because of an ongoing increased risk of meningococcal disease (e.g., those with complement deficiency, HIV, or asplenia): Follow the booster schedule for persons at increased risk (see below).
 - **Children for whom boosters are not recommended** (e.g., those who received a single dose for travel to a country where meningococcal disease is endemic): Administer MenACWY according to the recommended adolescent schedule with dose 1 at age 11–12 years and dose 2 at age 16 years.
- Note:** **Menactra** should be administered either before or at the same time as DTaP. For MenACWY **booster dose recommendations** for groups listed under “Special situations” and in an outbreak setting and for additional meningococcal vaccination information, see www.cdc.gov/vaccines/hcp/acip-recs/vacc-specific/mening.html.

Meningococcal serogroup B vaccination (minimum age: 10 years [MenB-4C, Bexsero; MenB-FHbp, Trumenba])

Shared clinical decision-making

- **Adolescents not at increased risk** age 16–23 years (preferred age 16–18 years) based on shared clinical decision-making:
 - **Bexsero:** 2-dose series at least 1 month apart
 - **Trumenba:** 2-dose series at least 6 months apart; if dose 2 is administered earlier than 6 months, administer a 3rd dose at least 4 months after dose 2.

Special situations

Anatomic or functional asplenia (including sickle cell disease), persistent complement component deficiency, complement inhibitor (e.g., eculizumab, ravulizumab) use:

- **Bexsero:** 2-dose series at least 1 month apart
- **Trumenba:** 3-dose series at 0, 1–2, 6 months

Bexsero and **Trumenba** are not interchangeable; the same product should be used for all doses in a series. For MenB **booster dose recommendations** for groups listed under “Special situations” and in an outbreak setting and for additional meningococcal vaccination information, see www.cdc.gov/vaccines/acip/recommendations.html and www.cdc.gov/vaccines/hcp/acip-recs/vacc-specific/mening.html.

Pneumococcal vaccination (minimum age: 6 weeks [PCV13], 2 years [PPSV23])

Routine vaccination with PCV13

- 4-dose series at 2, 4, 6, 12–15 months

Catch-up vaccination with PCV13

- 1 dose for healthy children age 24–59 months with any incomplete* PCV13 series
- For other catch-up guidance, see Table 2.

Special situations

High-risk conditions below: When both PCV13 and PPSV23 are indicated, administer PCV13 first. PCV13 and PPSV23 should not be administered during the same visit.

Chronic heart disease (particularly cyanotic congenital heart disease and cardiac failure), chronic lung disease (including asthma treated with high-dose, oral corticosteroids), diabetes mellitus:

Age 2–5 years

- Any incomplete* series with:
 - 3 PCV13 doses: 1 dose PCV13 (at least 8 weeks after any prior PCV13 dose)
 - Less than 3 PCV13 doses: 2 doses PCV13 (8 weeks after the most recent dose and administered 8 weeks apart)
- No history of PPSV23: 1 dose PPSV23 (at least 8 weeks after any prior PCV13 dose)

Age 6–18 years

- No history of PPSV23: 1 dose PPSV23 (at least 8 weeks after any prior PCV13 dose)

Cerebrospinal fluid leak, cochlear implant:

Age 2–5 years

- Any incomplete* series with:
 - 3 PCV13 doses: 1 dose PCV13 (at least 8 weeks after any prior PCV13 dose)
 - Less than 3 PCV13 doses: 2 doses PCV13 (8 weeks after the most recent dose and administered 8 weeks apart)
- No history of PPSV23: 1 dose PPSV23 (at least 8 weeks after any prior PCV13 dose)

Age 6–18 years

- No history of either PCV13 or PPSV23: 1 dose PCV13, 1 dose PPSV23 at least 8 weeks later
- Any PCV13 but no PPSV23: 1 dose PPSV23 at least 8 weeks after the most recent dose of PCV13
- PPSV23 but no PCV13: 1 dose PCV13 at least 8 weeks after the most recent dose of PPSV23

Sickle cell disease and other hemoglobinopathies; anatomic or functional asplenia; congenital or acquired immunodeficiency; HIV infection; chronic renal failure; nephrotic syndrome; malignant neoplasms, leukemias, lymphomas, Hodgkin disease, and other diseases associated with treatment with immunosuppressive drugs or radiation therapy; solid organ transplantation; multiple myeloma:

Age 2–5 years

- Any incomplete* series with:
 - 3 PCV13 doses: 1 dose PCV13 (at least 8 weeks after any prior PCV13 dose)
 - Less than 3 PCV13 doses: 2 doses PCV13 (8 weeks after the most recent dose and administered 8 weeks apart)
- No history of PPSV23: 1 dose PPSV23 (at least 8 weeks after any prior PCV13 dose) and a 2nd dose of PPSV23 5 years later

Age 6–18 years

- No history of either PCV13 or PPSV23: 1 dose PCV13, 2 doses PPSV23 (dose 1 of PPSV23 administered 8 weeks after PCV13 and dose 2 of PPSV23 administered at least 5 years after dose 1 of PPSV23)
- Any PCV13 but no PPSV23: 2 doses PPSV23 (dose 1 of PPSV23 administered 8 weeks after the most recent dose of PCV13 and dose 2 of PPSV23 administered at least 5 years after dose 1 of PPSV23)
- PPSV23 but no PCV13: 1 dose PCV13 at least 8 weeks after the most recent PPSV23 dose and a 2nd dose of PPSV23 administered 5 years after dose 1 of PPSV23 and at least 8 weeks after a dose of PCV13

Chronic liver disease, alcoholism:

Age 6–18 years

- No history of PPSV23: 1 dose PPSV23 (at least 8 weeks after any prior PCV13 dose)

**Incomplete series* = Not having received all doses in either the recommended series or an age-appropriate catch-up series. See Tables 8, 9, and 11 in the ACIP pneumococcal vaccine recommendations at www.cdc.gov/mmwr/pdf/rr/rr5911.pdf for complete schedule details.

Poliovirus vaccination (minimum age: 6 weeks)

Routine vaccination

- 4-dose series at ages 2, 4, 6–18 months, 4–6 years; administer the final dose at or after age 4 years and at least 6 months after the previous dose.
- 4 or more doses of IPV can be administered before age 4 years when a combination vaccine containing IPV is used. However, a dose is still recommended at or after age 4 years and at least 6 months after the previous dose.

Catch-up vaccination

- In the first 6 months of life, use minimum ages and intervals only for travel to a polio-endemic region or during an outbreak.
- IPV is not routinely recommended for U.S. residents 18 years and older.

Series containing oral polio vaccine (OPV), either mixed OPV-IPV or OPV-only series:

- Total number of doses needed to complete the series is the same as that recommended for the U.S. IPV schedule. See www.cdc.gov/mmwr/volumes/66/wr/mm6601a6.htm?s_cid=mm6601a6_w.
- Only trivalent OPV (tOPV) counts toward the U.S. vaccination requirements.
 - Doses of OPV administered before April 1, 2016, should be counted (unless specifically noted as administered during a campaign).
 - Doses of OPV administered on or after April 1, 2016, should not be counted.
 - For guidance to assess doses documented as “OPV”; see www.cdc.gov/mmwr/volumes/66/wr/mm6606a7.htm?s_cid=mm6606a7_w.
- For other catch-up guidance, see Table 2.

Rotavirus vaccination (minimum age: 6 weeks)

Routine vaccination

- **Rotarix:** 2-dose series at 2 and 4 months
- **RotaTeq:** 3-dose series at 2, 4, and 6 months
- If any dose in the series is either **RotaTeq** or unknown, default to 3-dose series.

Catch-up vaccination

- Do not start the series on or after age 15 weeks, 0 days.
- The maximum age for the final dose is 8 months, 0 days.
- For other catch-up guidance, see Table 2.

Tetanus, diphtheria, and pertussis (Tdap) vaccination

(minimum age: 11 years for routine vaccination, 7 years for catch-up vaccination)

Routine vaccination

- **Adolescents age 11–12 years:** 1 dose Tdap
- **Pregnancy:** 1 dose Tdap during each pregnancy, preferably in early part of gestational weeks 27–36
- Tdap may be administered regardless of the interval since the last tetanus- and diphtheria-toxoid-containing vaccine.

Catch-up vaccination

- **Adolescents age 13–18 years who have not received Tdap:** 1 dose Tdap, then Td or Tdap booster every 10 years
- **Persons age 7–18 years not fully vaccinated* with DTaP:** 1 dose Tdap as part of the catch-up series (preferably the first dose); if additional doses are needed, use Td or Tdap.
- **Tdap administered at 7–10 years:**
 - **Children age 7–9 years** who receive Tdap should receive the routine Tdap dose at age 11–12 years.
 - **Children age 10 years** who receive Tdap do not need to receive the routine Tdap dose at age 11–12 years.
- **DTaP inadvertently administered at or after age 7 years:**
 - **Children age 7–9 years:** DTaP may count as part of catch-up series. Routine Tdap dose at age 11–12 years should be administered.
 - **Children age 10–18 years:** Count dose of DTaP as the adolescent Tdap booster.
- For other catch-up guidance, see Table 2.
- For information on use of Tdap or Td as tetanus prophylaxis in wound management, see www.cdc.gov/mmwr/volumes/67/rr/rr6702a1.htm.

**Fully vaccinated* = 5 valid doses of DTaP OR 4 valid doses of DTaP if dose 4 was administered at age 4 years or older

Varicella vaccination (minimum age: 12 months)

Routine vaccination

- 2-dose series at 12–15 months, 4–6 years
- Dose 2 may be administered as early as 3 months after dose 1 (a dose administered after a 4-week interval may be counted).

Catch-up vaccination

- Ensure persons age 7–18 years without evidence of immunity (see www.cdc.gov/mmwr/pdf/rr/rr5604.pdf) have 2-dose series:
 - **Age 7–12 years:** routine interval: 3 months (a dose administered after a 4-week interval may be counted)
 - **Age 13 years and older:** routine interval: 4–8 weeks (minimum interval: 4 weeks)
 - The maximum age for use of MMRV is 12 years.

Recommended Adult Immunization Schedule for ages 19 years or older

UNITED STATES
2020

How to use the adult immunization schedule

- 1** Determine recommended vaccinations by age (**Table 1**)
- 2** Assess need for additional recommended vaccinations by medical condition and other indications (**Table 2**)
- 3** Review vaccine types, frequencies, and intervals and considerations for special situations (**Notes**)

Recommended by the Advisory Committee on Immunization Practices (www.cdc.gov/vaccines/acip) and approved by the Centers for Disease Control and Prevention (www.cdc.gov), American College of Physicians (www.acponline.org), American Academy of Family Physicians (www.aafp.org), American College of Obstetricians and Gynecologists (www.acog.org), and American College of Nurse-Midwives (www.midwife.org).

Vaccines in the Adult Immunization Schedule*

Vaccines	Abbreviations	Trade names
<i>Haemophilus influenzae</i> type b vaccine	Hib	ActHIB® Hiberix® PedvaxHIB®
Hepatitis A vaccine	HepA	Havrix® Vaqta®
Hepatitis A and hepatitis B vaccine	HepA-HepB	Twinrix®
Hepatitis B vaccine	HepB	Engerix-B® Recombivax HB® Heplisav-B®
Human papillomavirus vaccine	HPV vaccine	Gardasil 9®
Influenza vaccine (inactivated)	IIV	Many brands
Influenza vaccine (live, attenuated)	LAIV	FluMist® Quadrivalent
Influenza vaccine (recombinant)	RIV	Flublok® Quadrivalent
Measles, mumps, and rubella vaccine	MMR	M-M-R® II
Meningococcal serogroups A, C, W, Y vaccine	MenACWY	Menactra® Menveo®
Meningococcal serogroup B vaccine	MenB-4C MenB-FHbp	Bexsero® Trumenba®
Pneumococcal 13-valent conjugate vaccine	PCV13	Prenvar 13®
Pneumococcal 23-valent polysaccharide vaccine	PPSV23	Pneumovax® 23
Tetanus and diphtheria toxoids	Td	Tenivac® Tdvax™
Tetanus and diphtheria toxoids and acellular pertussis vaccine	Tdap	Adacel® Boostrix®
Varicella vaccine	VAR	Varivax®
Zoster vaccine, recombinant	RZV	Shingrix
Zoster vaccine live	ZVL	Zostavax®

*Administer recommended vaccines if vaccination history is incomplete or unknown. Do not restart or add doses to vaccine series if there are extended intervals between doses. The use of trade names is for identification purposes only and does not imply endorsement by the ACIP or CDC.

Report

- Suspected cases of reportable vaccine-preventable diseases or outbreaks to the local or state health department
- Clinically significant postvaccination reactions to the Vaccine Adverse Event Reporting System at www.vaers.hhs.gov or 800-822-7967

Injury claims

All vaccines included in the adult immunization schedule except pneumococcal 23-valent polysaccharide (PPSV23) and zoster (RZV, ZVL) vaccines are covered by the Vaccine Injury Compensation Program. Information on how to file a vaccine injury claim is available at www.hrsa.gov/vaccinecompensation.

Questions or comments

Contact www.cdc.gov/cdc-info or 800-CDC-INFO (800-232-4636), in English or Spanish, 8 a.m.–8 p.m. ET, Monday through Friday, excluding holidays.



Download the CDC Vaccine Schedules App for providers at www.cdc.gov/vaccines/schedules/hcp/schedule-app.html.

Helpful information

- Complete ACIP recommendations: www.cdc.gov/vaccines/hcp/acip-recs/index.html
- General Best Practice Guidelines for Immunization (including contraindications and precautions): www.cdc.gov/vaccines/hcp/acip-recs/general-recs/index.html
- Vaccine information statements: www.cdc.gov/vaccines/hcp/vis/index.html
- Manual for the Surveillance of Vaccine-Preventable Diseases (including case identification and outbreak response): www.cdc.gov/vaccines/pubs/surv-manual
- Travel vaccine recommendations: www.cdc.gov/travel
- Recommended Child and Adolescent Immunization Schedule, United States, 2020: www.cdc.gov/vaccines/schedules/hcp/child-adolescent.html



**U.S. Department of
Health and Human Services**
Centers for Disease
Control and Prevention

Table 1 Recommended Adult Immunization Schedule by Age Group, United States, 2020

Vaccine	19–26 years	27–49 years	50–64 years	≥65 years
Influenza inactivated (IIV) or Influenza recombinant (RIV)	1 dose annually			
Influenza live, attenuated (LAIV)	1 dose annually			
Tetanus, diphtheria, pertussis (Tdap or Td)	1 dose Tdap, then Td or Tdap booster every 10 years			
Measles, mumps, rubella (MMR)	1 or 2 doses depending on indication (if born in 1957 or later)			
Varicella (VAR)	2 doses (if born in 1980 or later)		2 doses	
Zoster recombinant (RZV) (preferred)			2 doses	
Zoster live (ZVL)			1 dose	
Human papillomavirus (HPV)	2 or 3 doses depending on age at initial vaccination or condition	27 through 45 years		
Pneumococcal conjugate (PCV13)	1 dose			65 years and older
Pneumococcal polysaccharide (PPSV23)	1 or 2 doses depending on indication			1 dose
Hepatitis A (HepA)	2 or 3 doses depending on vaccine			
Hepatitis B (HepB)	2 or 3 doses depending on vaccine			
Meningococcal A, C, W, Y (MenACWY)	1 or 2 doses depending on indication, see notes for booster recommendations			
Meningococcal B (MenB)	2 or 3 doses depending on vaccine and indication, see notes for booster recommendations			
	19 through 23 years			
Haemophilus influenzae type b (Hib)	1 or 3 doses depending on indication			

Recommended vaccination for adults who meet age requirement, lack documentation of vaccination, or lack evidence of past infection

Recommended vaccination for adults with an additional risk factor or another indication

Recommended vaccination based on shared clinical decision-making

No recommendation/ Not applicable

Table 2 Recommended Adult Immunization Schedule by Medical Condition and Other Indications, United States, 2020

Vaccine	Pregnancy	Immuno-compromised (excluding HIV infection)	HIV infection CD4 count		Asplenia, complement deficiencies	End-stage renal disease; or on hemodialysis	Heart or lung disease, alcoholism ¹	Chronic liver disease	Diabetes	Health care personnel ²	Men who have sex with men	
			<200	≥200								
IIV or RIV	1 dose annually											
LAIV	NOT RECOMMENDED					PRECAUTION				1 dose annually		
Tdap or Td	1 dose Tdap each pregnancy	1 dose Tdap, then Td or Tdap booster every 10 years										
MMR	NOT RECOMMENDED			1 or 2 doses depending on indication								
VAR	NOT RECOMMENDED			2 doses								
RZV (preferred)	DELAY				2 doses at age ≥50 years							
ZVL	NOT RECOMMENDED				1 dose at age ≥60 years							
HPV	DELAY	3 doses through age 26 years			2 or 3 doses through age 26 years							
PCV13		1 dose										
PPSV23		1, 2, or 3 doses depending on age and indication										
HepA							2 or 3 doses depending on vaccine					
HepB							2 or 3 doses depending on vaccine					
MenACWY	1 or 2 doses depending on indication, see notes for booster recommendations											
MenB	PRECAUTION	2 or 3 doses depending on vaccine and indication, see notes for booster recommendations										
Hib		3 doses HSCT ³ recipients only			1 dose							

Recommended vaccination for adults who meet age requirement, lack documentation of vaccination, or lack evidence of past infection
 Recommended vaccination for adults with an additional risk factor or another indication
 Precaution—vaccination might be indicated if benefit of protection outweighs risk of adverse reaction
 Delay vaccination until after pregnancy if vaccine is indicated
 Not recommended/contraindicated—vaccine should not be administered
 No recommendation/Not applicable

1. Precaution for LAIV does not apply to alcoholism. 2. See notes for influenza; hepatitis B; measles, mumps, and rubella; and varicella vaccinations. 3. Hematopoietic stem cell transplant.

Haemophilus influenzae* type b vaccination*Special situations**

- **Anatomical or functional asplenia (including sickle cell disease):** 1 dose if previously did not receive Hib; if elective splenectomy, 1 dose, preferably at least 14 days before splenectomy
- **Hematopoietic stem cell transplant (HSCT):** 3-dose series 4 weeks apart starting 6–12 months after successful transplant, regardless of Hib vaccination history

Hepatitis A vaccination**Routine vaccination**

- **Not at risk but want protection from hepatitis A** (identification of risk factor not required): 2-dose series HepA (Havrix 6–12 months apart or Vaqta 6–18 months apart [minimum interval: 6 months]) or 3-dose series HepA-HepB (Twinrix at 0, 1, 6 months [minimum intervals: 4 weeks between doses 1 and 2/5 months between doses 2 and 3])

Special situations

- **At risk for hepatitis A virus infection:** 2-dose series HepA or 3-dose series HepA-HepB as above
 - **Chronic liver disease** (e.g., persons with hepatitis B, hepatitis C, cirrhosis, fatty liver disease, alcoholic liver disease, autoimmune hepatitis, alanine aminotransferase [ALT] or aspartate aminotransferase [AST] level greater than twice the upper limit of normal)
 - **HIV infection**
 - **Men who have sex with men**
 - **Injection or noninjection drug use**
 - **Persons experiencing homelessness**
 - **Work with hepatitis A virus** in research laboratory or with nonhuman primates with hepatitis A virus infection
 - **Travel in countries with high or intermediate endemic hepatitis A**
 - **Close, personal contact with international adoptee** (e.g., household or regular babysitting) in first 60 days after arrival from country with high or intermediate endemic hepatitis A (administer dose 1 as soon as adoption is planned, at least 2 weeks before adoptee's arrival)

- **Pregnancy** if at risk for infection or severe outcome from infection during pregnancy
- **Settings for exposure, including** health care settings targeting services to injection or noninjection drug users or group homes and nonresidential day care facilities for developmentally disabled persons (individual risk factor screening not required)

Hepatitis B vaccination**Routine vaccination**

- **Not at risk but want protection from hepatitis B** (identification of risk factor not required): 2- or 3-dose series (2-dose series HepB at least 4 weeks apart [2-dose series HepB only applies when 2 doses of HepB are used at least 4 weeks apart] or 3-dose series Engerix-B or Recombivax HB at 0, 1, 6 months [minimum intervals: 4 weeks between doses 1 and 2/8 weeks between doses 2 and 3/16 weeks between doses 1 and 3]) or 3-dose series HepA-HepB (Twinrix at 0, 1, 6 months [minimum intervals: 4 weeks between doses 1 and 2/5 months between doses 2 and 3])

Special situations

- **At risk for hepatitis B virus infection:** 2-dose (HepB) or 3-dose (Engerix-B, Recombivax HB) series or 3-dose series HepA-HepB (Twinrix) as above
 - **Chronic liver disease** (e.g., persons with hepatitis C, cirrhosis, fatty liver disease, alcoholic liver disease, autoimmune hepatitis, alanine aminotransferase [ALT] or aspartate aminotransferase [AST] level greater than twice upper limit of normal)
 - **HIV infection**
 - **Sexual exposure risk** (e.g., sex partners of hepatitis B surface antigen [HBsAg]-positive persons; sexually active persons not in mutually monogamous relationships; persons seeking evaluation or treatment for a sexually transmitted infection; men who have sex with men)
 - **Current or recent injection drug use**
 - **Percutaneous or mucosal risk for exposure to blood** (e.g., household contacts of HBsAg-positive persons; residents and staff of facilities for developmentally disabled persons; health care and public safety personnel with reasonably anticipated risk for

exposure to blood or blood-contaminated body fluids; hemodialysis, peritoneal dialysis, home dialysis, and predialysis patients; persons with diabetes mellitus age younger than 60 years and, at discretion of treating clinician, those age 60 years or older)

- **Incarcerated persons**
- **Travel in countries with high or intermediate endemic hepatitis B**

- **Pregnancy** if at risk for infection or severe outcome from infection during pregnancy (HepB not currently recommended due to lack of safety data in pregnant women)

Human papillomavirus vaccination**Routine vaccination**

- **HPV vaccination recommended for all adults through age 26 years:** 2- or 3-dose series depending on age at initial vaccination or condition:
 - **Age 15 years or older at initial vaccination:** 3-dose series at 0, 1–2, 6 months (minimum intervals: 4 weeks between doses 1 and 2/12 weeks between doses 2 and 3/5 months between doses 1 and 3; repeat dose if administered too soon)
 - **Age 9 through 14 years at initial vaccination and received 1 dose or 2 doses less than 5 months apart:** 1 dose
 - **Age 9 through 14 years at initial vaccination and received 2 doses at least 5 months apart:** HPV vaccination complete, no additional dose needed.
- **If completed valid vaccination series with any HPV vaccine, no additional doses needed**

Shared clinical decision-making

- **Age 27 through 45 years based on shared clinical decision-making:**
 - 2- or 3-dose series as above

Special situations

- **Pregnancy through age 26 years:** HPV vaccination is not recommended until after pregnancy; no intervention needed if vaccinated while pregnant; pregnancy testing not needed before vaccination

Influenza vaccination

Routine vaccination

- **Persons age 6 months or older:** 1 dose any influenza vaccine appropriate for age and health status annually
- For additional guidance, see www.cdc.gov/flu/professionals/index.htm

Special situations

- **Egg allergy, hives only:** 1 dose any influenza vaccine appropriate for age and health status annually
- **Egg allergy more severe than hives** (e.g., angioedema, respiratory distress): 1 dose any influenza vaccine appropriate for age and health status annually in medical setting under supervision of health care provider who can recognize and manage severe allergic reactions
- **LAIV should not be used** in persons with the following conditions or situations:
 - History of severe allergic reaction to any vaccine component (excluding egg) or to a previous dose of any influenza vaccine
 - Immunocompromised due to any cause (including medications and HIV infection)
 - Anatomic or functional asplenia
 - Cochlear implant
 - Cerebrospinal fluid-oro-pharyngeal communication
 - Close contacts or caregivers of severely immunosuppressed persons who require a protected environment
 - Pregnancy
 - Received influenza antiviral medications within the previous 48 hours
- **History of Guillain-Barré syndrome within 6 weeks of previous dose of influenza vaccine:** Generally should not be vaccinated unless vaccination benefits outweigh risks for those at higher risk for severe complications from influenza

Measles, mumps, and rubella vaccination

Routine vaccination

- **No evidence of immunity to measles, mumps, or rubella:** 1 dose
 - **Evidence of immunity:** Born before 1957 (health care personnel, see below), documentation of receipt of MMR vaccine, laboratory evidence of immunity or disease (diagnosis of disease without laboratory confirmation is not evidence of immunity)

Special situations

- **Pregnancy with no evidence of immunity to rubella:** MMR contraindicated during pregnancy; after pregnancy (before discharge from health care facility), 1 dose
- **Nonpregnant women of childbearing age with no evidence of immunity to rubella:** 1 dose
- **HIV infection with CD4 count ≥ 200 cells/ μ L for at least 6 months and no evidence of immunity to measles, mumps, or rubella:** 2-dose series at least 4 weeks apart; MMR contraindicated in HIV infection with CD4 count < 200 cells/ μ L
- **Severe immunocompromising conditions:** MMR contraindicated
- **Students in postsecondary educational institutions, international travelers, and household or close, personal contacts of immunocompromised persons with no evidence of immunity to measles, mumps, or rubella:** 2-dose series at least 4 weeks apart if previously did not receive any doses of MMR or 1 dose if previously received 1 dose MMR
- **Health care personnel:**
 - **Born in 1957 or later with no evidence of immunity to measles, mumps, or rubella:** 2-dose series at least 4 weeks apart for measles or mumps or at least 1 dose for rubella
 - **Born before 1957 with no evidence of immunity to measles, mumps, or rubella:** Consider 2-dose series at least 4 weeks apart for measles or mumps or 1 dose for rubella

Meningococcal vaccination

Special situations for MenACWY

- **Anatomical or functional asplenia (including sickle cell disease), HIV infection, persistent complement component deficiency, complement inhibitor (e.g., eculizumab, ravulizumab) use:** 2-dose series MenACWY (Menactra, Menveo) at least 8 weeks apart and revaccinate every 5 years if risk remains
- **Travel in countries with hyperendemic or epidemic meningococcal disease, microbiologists routinely exposed to *Neisseria meningitidis*:** 1 dose MenACWY (Menactra, Menveo) and revaccinate every 5 years if risk remains
- **First-year college students who live in residential housing (if not previously vaccinated at age 16 years or older) and military recruits:** 1 dose MenACWY (Menactra, Menveo)

Shared clinical decision-making for MenB

- **Adolescents and young adults age 16 through 23 years (age 16 through 18 years preferred) not at increased risk for meningococcal disease:** Based on shared clinical decision-making, 2-dose series MenB-4C at least 1 month apart or 2-dose series MenB-FHbp at 0, 6 months (if dose 2 was administered less than 6 months after dose 1, administer dose 3 at least 4 months after dose 2); MenB-4C and MenB-FHbp are not interchangeable (use same product for all doses in series)

Special situations for MenB

- **Anatomical or functional asplenia (including sickle cell disease), persistent complement component deficiency, complement inhibitor (e.g., eculizumab, ravulizumab) use, microbiologists routinely exposed to *Neisseria meningitidis*:** 2-dose primary series MenB-4C (Bexsero) at least 1 month apart or 3-dose primary series MenB-FHbp (Trumenba) at 0, 1–2, 6 months (if dose 2 was administered at least 6 months after dose 1, dose 3 not needed); MenB-4C and MenB-FHbp are not interchangeable (use same product for all doses in series); 1 dose MenB booster 1 year after primary series and revaccinate every 2–3 years if risk remains
- **Pregnancy:** Delay MenB until after pregnancy unless at increased risk and vaccination benefits outweigh potential risks

Pneumococcal vaccination

Routine vaccination

- **Age 65 years or older** (immunocompetent—see www.cdc.gov/mmwr/volumes/68/wr/mm6846a5.htm?s_cid=mm6846a5_w): 1 dose PPSV23
 - If PPSV23 was administered prior to age 65 years, administer 1 dose PPSV23 at least 5 years after previous dose

Shared clinical decision-making

- **Age 65 years and older** (immunocompetent): 1 dose PCV13 based on **shared clinical decision-making**
 - If both PCV13 and PPSV23 are to be administered, PCV13 should be administered first
 - PCV13 and PPSV23 should be administered at least 1 year apart
 - PCV13 and PPSV23 should not be administered during the same visit

Special situations

(see www.cdc.gov/mmwr/volumes/68/wr/mm6846a5.htm?s_cid=mm6846a5_w)

- **Age 19 through 64 years with chronic medical conditions (chronic heart [excluding hypertension], lung, or liver disease, diabetes), alcoholism, or cigarette smoking:** 1 dose PPSV23
- **Age 19 years or older with immunocompromising conditions (congenital or acquired immunodeficiency [including B- and T-lymphocyte deficiency, complement deficiencies, phagocytic disorders, HIV infection], chronic renal failure, nephrotic syndrome, leukemia, lymphoma, Hodgkin disease, generalized malignancy, iatrogenic immunosuppression [e.g., drug or radiation therapy], solid organ transplant, multiple myeloma) or anatomical or functional asplenia (including sickle cell disease and other hemoglobinopathies):** 1 dose PCV13 followed by 1 dose PPSV23 at least 8 weeks later, then another dose PPSV23 at least 5 years after previous PPSV23; at age 65 years or older, administer 1 dose PPSV23 at least 5 years after most recent PPSV23 (note: only 1 dose PPSV23 recommended at age 65 years or older)

- **Age 19 years or older with cerebrospinal fluid leak or cochlear implant:** 1 dose PCV13 followed by 1 dose PPSV23 at least 8 weeks later; at age 65 years or older, administer another dose PPSV23 at least 5 years after PPSV23 (note: only 1 dose PPSV23 recommended at age 65 years or older)

Tetanus, diphtheria, and pertussis vaccination

Routine vaccination

- **Previously did not receive Tdap at or after age 11 years:** 1 dose Tdap, then Td or Tdap every 10 years

Special situations

- **Previously did not receive primary vaccination series for tetanus, diphtheria, or pertussis:** At least 1 dose Tdap followed by 1 dose Td or Tdap at least 4 weeks after Tdap and another dose Td or Tdap 6–12 months after last Td or Tdap (Tdap can be substituted for any Td dose, but preferred as first dose); Td or Tdap every 10 years thereafter
- **Pregnancy:** 1 dose Tdap during each pregnancy, preferably in early part of gestational weeks 27–36
- For information on use of Td or Tdap as tetanus prophylaxis in wound management, see www.cdc.gov/mmwr/volumes/67/rr/rr6702a1.htm

Varicella vaccination

Routine vaccination

- **No evidence of immunity to varicella:** 2-dose series 4–8 weeks apart if previously did not receive varicella-containing vaccine (VAR or MMRV [measles-mumps-rubella-varicella vaccine] for children); if previously received 1 dose varicella-containing vaccine, 1 dose at least 4 weeks after first dose
 - Evidence of immunity: U.S.-born before 1980 (except for pregnant women and health care personnel [see below]), documentation of 2 doses varicella-containing vaccine at least 4 weeks apart, diagnosis or verification of history of varicella or herpes zoster by a health care provider, laboratory evidence of immunity or disease

Special situations

- **Pregnancy with no evidence of immunity to varicella:** VAR contraindicated during pregnancy; after pregnancy (before discharge from health care facility) 1 dose if previously received 1 dose varicella-containing vaccine or dose 1 of 2-dose series (dose 2: 4–8 weeks later) if previously did not receive any varicella-containing vaccine, regardless of whether U.S.-born before 1980
- **Health care personnel with no evidence of immunity to varicella:** 1 dose if previously received 1 dose varicella-containing vaccine; 2-dose series 4–8 weeks apart if previously did not receive any varicella-containing vaccine, regardless of whether U.S.-born before 1980
- **HIV infection with CD4 count ≥ 200 cells/ μ L with no evidence of immunity:** Vaccination may be considered (2 doses, administered 3 months apart); VAR contraindicated in HIV infection with CD4 count < 200 cells/ μ L
- **Severe immunocompromising conditions:** VAR contraindicated

Zoster vaccination

Routine vaccination

- **Age 50 years or older:** 2-dose series RZV (Shingrix) 2–6 months apart (minimum interval: 4 weeks; repeat dose if administered too soon), regardless of previous herpes zoster or history of ZVL (Zostavax) vaccination (administer RZV at least 2 months after ZVL)
- **Age 60 years or older:** 2-dose series RZV 2–6 months apart (minimum interval: 4 weeks; repeat if administered too soon) or 1 dose ZVL if not previously vaccinated. RZV preferred over ZVL (if previously received ZVL, administer RZV at least 2 months after ZVL)

Special situations

- **Pregnancy:** ZVL contraindicated; consider delaying RZV until after pregnancy if RZV is otherwise indicated
- **Severe immunocompromising conditions (including HIV infection with CD4 count < 200 cells/ μ L):** ZVL contraindicated; recommended use of RZV under review

Attachment 4



[WORKFORCE INTRODUCTION HERE]



1. Are you currently licensed as a pharmacist in California?

- Yes
 No

2. Is your primary practice setting located in California?

- Yes
 No

3. Which of the following best describes your primary practice setting?

4. Are you the designated pharmacist-in-charge?

- Yes
 No

5. Are you in a management position for your employer, e.g., pharmacy manager, district manager?

- Yes
 No

6. Do you work at multiple worksites for a single employer or through a relief agency?

- Yes
 No



7. Please describe your typical work shift at your primary work site (e.g, day shift, swing shift, night shift, etc.)

8. Approximately how long is a typical shift at your primary work site?

- 8 hours
- 10 hours
- 12 hours
- Other (please specify)

9. Please provide the number of staff that work during a typical shift at your primary work site

Number of Pharmacists

Number of Interns

Number of Pharmacy
Technicians

Number of Unlicensed
Clerk/Typists

Not Applicable for my
primary worksite

10. What is the average prescription volume during a typical shift at your primary worksite?

- Less than 50
- 50 – 100 prescriptions
- 101 – 150 prescriptions
- 151 – 200 prescriptions
- 201 – 250 prescriptions
- 251 – 300 prescriptions
- Over 300 prescriptions
- Not Applicable for my primary worksite

11. Does your primary work-site perform sterile compounding?

- Yes
- No

If yes, please specify the average number of preparations compounded in a typical shift.

12. Does your primary work-site perform non-sterile compounding?

- Yes
- No

If yes, please specify the average number of preparations compounded in a typical shift.

13. Which of the following additional patient services are provided at your primary work site? (Please select all that apply.)

- Immunizations
- Hormonal contraception
- Travel medications
- Naloxone
- Smoking cessation
- CLIA waived tests
- Administer medication and biologicals
- Not Applicable for my primary worksite
- Other (please specify)

14. If your primary worksite provides immunizations, please indicate the average number administered during a typical work shift.

15. Does your primary work site use an automated drug delivery system such as Script Pro or Yuyama?

- Yes
- No
- Not Applicable for my primary worksite

If yes, (please specify type)

16. Does your primary worksite employer use workload metrics in the following areas? (Please select all that apply.)

- Number of immunizations administered.
- Number of refill requests
- Number of patients enrolled in an auto-refill program
- Number of prescriptions filled
- Number of M₁ completed
- Average time to fill a new prescription
- Average time to fill a refill prescription for a patient who is waiting
- No workload metrics are used
- Other (please specify)

17. Does your primary worksite have a work queue that monitors the dispensing time for a prescription?

- Yes
- No
- Not Applicable for my primary worksite

18. Does your employer at your primary worksite require other non-dispensing related duties (ringing up patients, making patient calls, developing staff schedules, meetings, etc.)?

- Yes
- No

Please detail the duties and the estimated percentage of your work shift dedicated to those duties.



19. On average, what is the number of medication errors that occur during a month at your primary worksite?

- Not Applicable for my primary worksite
 Insert Number

20. In your opinion, are medication errors appropriately documented and evaluated consistently with the Board's quality assurance requirements at your primary worksite?

- Yes
 No
 Not Applicable for my primary worksite

If no, please explain

21. Of the documented medication errors that occurred at your primary worksite, what were the cause for errors? (Please check all that apply.)

- Incorrect drug
 Incorrect strength
 Incorrect directions
 Incorrect prescriber
 Incorrect quantity
 Incorrect patient name
 Patient received prescription intended for another patient

22. Do you believe you have sufficient time to provide appropriate patient consultation?

- Yes
 No
 Not Applicable, my position does not require patient consultation

Additional Comments

23. Do you believe the pharmacy staffing in your primary worksite is appropriate to ensure adequate patient care?

Yes

No

24. Please provide any additional information you believe is important for the Board to understand about your worksite, that could impact patient care and medication errors.

Attachment 5

CALIFORNIA STATE BOARD OF PHARMACY
 QUARTERLY LICENSING STATISTICS FISCAL YEAR 2020/2021

APPLICATIONS RECEIVED

Individual Applications	July - Sept	Oct-Dec	Jan-Mar	Apr-Jun	Total FYTD
Designated Representatives (EXC)	92	90	0	0	182
Designated Representatives Vet (EXV)	0	1	0	0	1
Designated Representatives-3PL (DRL)	27	29	0	0	56
Designated Representatives-Reverse Distributor (DRR)	1	1	0	0	2
Designated Paramedic (DPM)	0	0	0	0	0
Intern Pharmacist (INT)	1,237	111	0	0	1,348
Pharmacist Exam Applications	299	152	0	0	451
Pharmacist Retake Exam Applications	585	489	0	0	1,074
Pharmacist Initial License Application (RPH)	935	555	0	0	1,490
Advanced Practice Pharmacist (APH)	60	44	0	0	104
Pharmacy Technician (TCH)	1,182	1,036	0	0	2,218
Total	4,418	2,508	0	0	6,926

Site Applications	July - Sept	Oct-Dec	Jan-Mar	Apr-Jun	Total FYTD
Automated Drug Delivery System (ADD)	50	60	0	0	110
Automated Drug Delivery System EMS (ADE)	0	0	0	0	0
Automated Patient Dispensing System 340B Clinic (ADC)	0	0	0	0	0
Centralized Hospital Packaging Government Owned (CHE)	0	0	0	0	0
Centralized Hospital Packaging (CHP)	1	0	0	0	1
Clinics (CLN)	26	40	0	0	66
Clinics Government Owned (CLE)	19	13	0	0	32
Drug Room (DRM)	0	0	0	0	0
Drug Room Government Owned (DRE)	0	0	0	0	0
Hospitals (HSP)	6	9	0	0	15
Hospitals Government Owned (HPE)	1	0	0	0	1
Hospital Satellite Sterile Compounding (SCP)	0	0	0	0	0
Hospital Satellite Sterile Compounding Government Owned (SCE)	0	0	0	0	0
Hypodermic Needle and Syringes (HYP)	4	5	0	0	9
Correctional Pharmacy (LCF)	0	0	0	0	0
Outsourcing Facility (OSF)	0	0	0	0	0
Outsourcing Facility Nonresident (NSF)	1	4	0	0	5
Pharmacy (PHY)	84	108	0	0	192
Pharmacy (PHY) Chain	5	8	0	0	13
Pharmacy Government Owned (PHE)	4	2	0	0	6
Remote Dispensing Pharmacy (PHR)	2	0	0	0	2
Pharmacy Nonresident (NRP)	30	33	0	0	63
Sterile Compounding (LSC)	23	37	0	0	60
Sterile Compounding Government Owned (LSE)	1	1	0	0	2
Sterile Compounding Nonresident (NSC)	5	2	0	0	7
Surplus Medication Collection Distribution Intermediary (SME)	0	0	0	0	0
Third-Party Logistics Providers (TPL)	4	2	0	0	6
Third-Party Logistics Providers Nonresident (NPL)	8	8	0	0	16
Veterinary Food-Animal Drug Retailer (VET)	0	0	0	0	0
Wholesalers (WLS)	20	9	0	0	29
Wholesalers Government Owned (WLE)	0	6	0	0	6
Wholesalers Nonresident (OSD)	30	26	0	0	56
Total	324	373	0	0	697

Applications Received with Temporary License Requests	July - Sept	Oct-Dec	Jan-Mar	Apr-Jun	Total FYTD
Drug Room -Temp (DRM)	0	2	0	0	2
Hospitals - Temp (HSP)	6	10	0	0	16
Hospital Satellite Sterile Compounding - Temp (SCP)	0	0	0	0	0
Outsourcing Facility - Temp (OSF)	0	0	0	0	0
Outsourcing Facility Nonresident - Temp (NSF)	0	0	0	0	0
Pharmacy - Temp (PHY)	58	77	0	0	135
Remote Dispensing Pharmacy - Temp (PHR)	1	0	0	0	1
Pharmacy Nonresident - Temp (NRP)	18	21	0	0	39
Sterile Compounding - Temp (LSC)	17	21	0	0	38
Sterile Compounding Nonresident - Temp (NSC)	1	1	0	0	2
Third-Party Logistics Providers - Temp (TPL)	3	0	0	0	3
Third-Party Logistics Providers Nonresident - Temp (NPL)	7	6	0	0	13
Veterinary Food-Animal Drug Retailer - Temp (VET)	0	0	0	0	0
Wholesalers - Temp (WLS)	6	6	0	0	12
Wholesalers Nonresident - Temp (OSD)	13	1	0	0	14
Total	130	145	0	0	275

LICENSES ISSUED

Individual Licenses	July - Sept	Oct-Dec	Jan-Mar	Apr-Jun	Total FYTD
Designated Representatives (EXC)	57	63	0	0	120
Designated Representatives Vet (EXV)	2	0	0	0	2
Designated Representatives-3PL (DRL)	18	19	0	0	37
Designated Representatives-Reverse Distributor (DRR)	0	1	0	0	1
Designated Paramedic (DPM)	0	0	0	0	0
Intern Pharmacist (INT)	935	389	0	0	1,324
Pharmacist (RPH)	936	557	0	0	1,493
Advanced Practice Pharmacist (APH)	34	41	0	0	75
Pharmacy Technician (TCH)	711	762	0	0	1,473
Total	2,693	1,832	0	0	4,525

Site Licenses	July - Sept	Oct-Dec	Jan-Mar	Apr-Jun	Total FYTD
Automated Drug Delivery System (ADD)	28	54	0	0	82
Automated Drug Delivery System EMS (ADE)	0	0	0	0	0
Automated Patient Dispensing System 340B Clinic (ADC)	0	0	0	0	0
Centralized Hospital Packaging Government Owned (CHE)	0	0	0	0	0
Centralized Hospital Packaging (CHP)	0	0	0	0	0
Clinics (CLN)	18	7	0	0	25
Clinics Government Owned (CLE)	18	13	0	0	31
Drug Room (DRM)	0	0	0	0	0
Drug Room Government Owned (DRE)	0	0	0	0	0
Hospitals (HSP)	0	0	0	0	0
Hospitals Government Owned (HPE)	0	0	0	0	0
Hospital Satellite Sterile Compounding (SCP)	0	0	0	0	0
Hospital Satellite Sterile Compounding Government Owned (SCE)	1	0	0	0	1
Hypodermic Needle and Syringes (HYP)	1	0	0	0	1
Correctional Pharmacy (LCF)	0	0	0	0	0
Outsourcing Facility (OSF)	0	0	0	0	0
Outsourcing Facility Nonresident (NSF)	1	0	0	0	1
Pharmacy (PHY)	27	15	0	0	42
Pharmacy Government Owned (PHE)	1	1	0	0	2
Remote Dispensing Pharmacy (PHR)	2	0	0	0	1
Pharmacy Nonresident (NRP)	4	0	0	0	4
Sterile Compounding (LSC)	10	7	0	0	17
Sterile Compounding Government Owned (LSE)	1	0	0	0	1
Sterile Compounding Nonresident (NSC)	0	0	0	0	0
Surplus Medication Collection Distribution Intermediary (SME)	0	0	0	0	0
Third-Party Logistics Providers (TPL)	0	1	0	0	1
Third-Party Logistics Providers Nonresident (NPL)	1	2	0	0	3
Veterinary Food-Animal Drug Retailer (VET)	0	0	0	0	0
Wholesalers (WLS)	5	5	0	0	10
Wholesalers Government Owned (WLE)	0	0	0	0	0
Wholesalers Nonresident (OSD)	15	4	0	0	19
Total	133	109	0	0	241

Site Temporary Licenses	July - Sept	Oct-Dec	Jan-Mar	Apr-Jun	Total FYTD
Drug Room -Temp (DRM)	0	1	0	0	1
Hospitals - Temp (HSP)	4	6	0	0	10
Hospital Satellite Sterile Compounding - Temp (SCP)	0	0	0	0	0
Outsourcing Facility - Temp (OSF)	1	0	0	0	1
Outsourcing Facility Nonresident - Temp (NSF)	1	0	0	0	1
Pharmacy - Temp (PHY)	49	49	0	0	98
Remote Dispensing Pharmacy - Temp (PHR)	2	0	0	0	2
Pharmacy Nonresident - Temp (NRP)	17	9	0	0	26
Sterile Compounding - Temp (LSC)	11	5	0	0	16
Sterile Compounding Nonresident - Temp (NSC)	1	0	0	0	1
Third-Party Logistics Providers - Temp (TPL)	0	2	0	0	2
Third-Party Logistics Providers Nonresident - Temp (NPL)	1	7	0	0	8
Veterinary Food-Animal Drug Retailer - Temp (VET)	0	0	0	0	0
Wholesalers - Temp (WLS)	4	8	0	0	12
Wholesalers Nonresident - Temp (OSD)	11	4	0	0	15
Total	102	91	0	0	193

PENDING APPLICATIONS (Data reflects number of pending applications at the end of the quarter)

Individual Applications	July - Sept	Oct-Dec	Jan-Mar	Apr-Jun
Designated Representatives (EXC)	178	206		
Designated Representatives (EXV)	2	3		
Designated Representatives-3PL (DRL)	42	54		
Designated Representatives-Reverse Distributor (DRR)	2	2		
Designated Paramedic (DPM)	0	0		
Intern Pharmacist (INT)	410	125		
Pharmacist (exam not eligible)	1,343	1,351		
Pharmacist (exam eligible)	1,456	941		
Advanced Practice Pharmacist (APH)	85	85		
Pharmacy Technician (TCH)	1,499	1,472		
Total	5,017	4,239	0	0

Site Applications	July - Sept	Oct-Dec	Jan-Mar	Apr-Jun
Automated Drug Delivery System (ADD)	163	179		
Automated Drug Delivery System EMS (ADE)	0	0		
Automated Patient Dispensing System 340B Clinic (ADC)	0	0		
Centralized Hospital Packaging Government Owned (CHE)	1	1		
Centralized Hospital Packaging (CHP)	5	5		
Clinics (CLN)	95	116		
Clinics Government Owned (CLE)	29	29		
Drug Room (DRM)	1	1		
Drug Room Government Owned (DRE)	0	0		
Hospitals (HSP)	19	16		
Hospitals Government Owned (HPE)	3	2		
Hospital Satellite Sterile Compounding (SCP)	2	2		
Hospital Satellite Sterile Compounding Government Owned (SCE)	0	0		
Hypodermic Needle and Syringes (HYP)	5	10		
Correctional Pharmacy (LCF)	0	0		
Outsourcing Facility (OSF)	0	0		
Outsourcing Facility Nonresident (NSF)	5	9		
Pharmacy (PHY)	150	190		
Pharmacy Government Owned (PHE)	4	5		
Remote Dispensing Pharmacy (PHR)	3	3		
Pharmacy Nonresident (NRP)	132	155		
Sterile Compounding (LSC)	84	106		
Sterile Compounding - Government Owned (LSE)	10	11		
Sterile Compounding Nonresident (NSC)	11	12		
Surplus Medication Collection Distribution Intermediary (SME)	0	0		
Third-Party Logistics Providers (TPL)	4	3		
Third-Party Logistics Providers Nonresident (NPL)	46	46		
Veterinary Food-Animal Drug Retailer (VET)	1	1		
Wholesalers (WLS)	46	50		
Wholesalers Government Owned (WLE)	1	1		
Wholesalers Nonresident (OSD)	91	109		
Total	911	1,062	0	0

Applications Pending with Temporary Licenses Issued - Pending Full License	July - Sept	Oct-Dec	Jan-Mar	Apr-Jun
Drug Room -Temp (DRM)	0	1		
Hospitals - Temp (HSP)	4	15		
Hospital Satellite Sterile Compounding - Temp (SCP)	0	0		
Outsourcing Facility - Temp (OSF)	1	0		
Outsourcing Facility Nonresident - Temp (NSF)	3	3		
Pharmacy - Temp (PHY)	99	91		
Remote Dispensing Pharmacy - Temp (PHR)	2	2		
Pharmacy Nonresident - Temp (NRP)	36	25		
Sterile Compounding - Temp (LSC)	11	13		
Sterile Compounding Nonresident - Temp (NSC)	2	1		
Third-Party Logistics Providers - Temp (TPL)	1	0		
Third-Party Logistics Providers Nonresident - Temp (NPL)	0	1		
Veterinary Food-Animal Drug Retailer - Temp (VET)	0	0		
Wholesalers - Temp (WLS)	4	9		
Wholesalers Nonresident - Temp (OSD)	9	9		
Total	172	170	0	0

APPLICATIONS WITHDRAWN

Individual Applications	July - Sept	Oct-Dec	Jan-Mar	Apr-Jun	Total FYTD
Designated Representatives (EXC)	233	4	0	0	237
Designated Representatives Vet (EXV)	1	0	0	0	1
Designated Representatives-3PL (DRL)	70	0	0	0	70
Designated Representatives-Reverse Distributor (DRR)	1	0	0	0	1
Designated Paramedic (DPM)	0	0	0	0	0
Intern Pharmacist (INT)	3	5	0	0	8
Pharmacist (exam applications)	239	430	0	0	669
Advanced Practice Pharmacist (APH)	9	4	0	0	13
Pharmacy Technician (TCH)	7	5	0	0	12
Total	563	448	0	0	1,011

Site Applications	July - Sept	Oct-Dec	Jan-Mar	Apr-Jun	Total FYTD
Automated Drug Delivery System (ADD)	3	2	0	0	5
Automated Drug Delivery System EMS (ADE)	0	0	0	0	0
Automated Patient Dispensing System 340B Clinic (ADC)	0	0	0	0	0
Centralized Hospital Packaging Government Owned (CHE)	0	0	0	0	0
Centralized Hospital Packaging (CHP)	0	0	0	0	0
Clinics (CLN)	1	8	0	0	9
Clinics Government Owned (CLE)	0	0	0	0	0
Drug Room (DRM)	0	0	0	0	0
Drug Room Government Owned (DRE)	0	0	0	0	0
Hospitals (HSP)	0	0	0	0	0
Hospitals Government Owned (HPE)	0	0	0	0	0
Hospital Satellite Sterile Compounding (SCP)	0	0	0	0	0
Hospital Satellite Sterile Compounding Government Owned (SCE)	1	0	0	0	1
Hypodermic Needle and Syringes (HYP)	0	0	0	0	0
Correctional Pharmacy (LCF)	0	0	0	0	0
Outsourcing Facility (OSF)	0	0	0	0	0
Outsourcing Facility Nonresident (NSF)	0	0	0	0	0
Pharmacy (PHY)	9	7	0	0	16
Pharmacy Government Owned (PHE)	2	0	0	0	2
Remote Dispensing Pharmacy (PHR)	0	0	0	0	0
Pharmacy Nonresident (NRP)	0	1	0	0	1
Sterile Compounding (LSC)	1	1	0	0	2
Sterile Compounding - Government Owned (LSE)	0	0	0	0	0
Sterile Compounding Nonresident (NSC)	0	0	0	0	0
Surplus Medication Collection Distribution Intermediary (SME)	0	0	0	0	0
Third-Party Logistics Providers (TPL)	0	0	0	0	0
Third-Party Logistics Providers Nonresident (NPL)	3	0	0	0	3
Veterinary Food-Animal Drug Retailer (VET)	0	0	0	0	0
Wholesalers (WLS)	1	0	0	0	1
Wholesalers Government Owned (WLE)	0	0	0	0	0
Wholesalers Nonresident (OSD)	1	1	0	0	2
Total	22	20	0	0	42

APPLICATIONS DENIED

Individual Applications	July - Sept	Oct-Dec	Jan-Mar	Apr-Jun	Total FYTD
Designated Representatives (EXC)	0	0	0	0	0
Designated Representatives Vet (EXV)	0	0	0	0	0
Designated Representatives-3PL (DRL)	0	0	0	0	0
Designated Representatives-Reverse Distributor (DRR)	0	0	0	0	0
Designated Paramedic (DPM)	0	0	0	0	0
Intern Pharmacist (INT)	0	0	0	0	0
Pharmacist (exam application)	1	1	0	0	2
Pharmacist (exam eligible)	0	0	0	0	0
Advanced Practice Pharmacist (APH)	0	0	0	0	0
Pharmacy Technician (TCH)	1	2	0	0	3
Total	2	3	0	0	5

Site Applications	July - Sept	Oct-Dec	Jan-Mar	Apr-Jun	Total FYTD
Centralized Hospital Packaging Government Owned (CHE)	0	0	0	0	0
Centralized Hospital Packaging (CHP)	0	0	0	0	0
Clinics (CLN)	0	0	0	0	0
Clinics Government Owned (CLE)	0	0	0	0	0
Drug Room (DRM)	0	0	0	0	0
Drug Room Government Owned (DRE)	0	0	0	0	0
Hospitals (HSP)	0	0	0	0	0
Hospitals Government Owned (HPE)	0	0	0	0	0
Hospital Satellite Sterile Compounding (SCP)	0	0	0	0	0
Hospital Satellite Sterile Compounding Government Owned (SCE)	0	0	0	0	0
Hypodermic Needle and Syringes (HYP)	0	0	0	0	0
Correctional Pharmacy (LCF)	0	0	0	0	0
Outsourcing Facility (OSF)	0	0	0	0	0
Outsourcing Facility Nonresident (NSF)	0	0	0	0	0
Pharmacy (PHY)	2	4	0	0	6
Pharmacy Government Owned (PHE)	0	0	0	0	0
Remote Dispensing Pharmacy (PHR)	0	0	0	0	0
Pharmacy Nonresident (NRP)	3	1	0	0	4
Sterile Compounding (LSC)	0	0	0	0	0
Sterile Compounding Government Owned (LSE)	0	0	0	0	0
Sterile Compounding Nonresident (NSC)	1	1	0	0	2
Surplus Medication Collection Distribution Intermediary (SME)	0	0	0	0	0
Third-Party Logistics Providers (TPL)	0	0	0	0	0
Third-Party Logistics Providers Nonresident (NPL)	0	0	0	0	0
Veterinary Food-Animal Drug Retailer (VET)	0	0	0	0	0
Wholesalers (WLS)	0	0	0	0	0
Wholesalers Government Owned (WLE)	0	0	0	0	0
Wholesalers Nonresident (OSD)	0	0	0	0	0
Total	6	6	0	0	12

RESPOND TO STATUS INQUIRIES

Email Inquiries	July - Sept	Oct-Dec	Jan-Mar	Apr-Jun	Total FYTD
Designated Representative Received	1,123	364	0	0	1,487
Designated Representative Responded	1,018	165	0	0	1,183
Advanced Practice Pharmacist Received	110	203	0	0	313
Advanced Practice Pharmacist Responded	110	217	0	0	327
Pharmacist/Intern Received	2,473	1,876	0	0	4,349
Pharmacist/Intern Responded	1,796	1,773	0	0	3,569
Pharmacy Technician Received	1,215	2,058	0	0	3,273
Pharmacy Technician Responded	1,193	1,325	0	0	2,518
Pharmacy Received	2,013	1,272	0	0	3,285
Pharmacy Responded	1,799	1,561	0	0	3,360
Sterile Compounding/Outsourcing Received	1,196	1,276	0	0	2,472
Sterile Compounding/Outsourcing Responded	630	846	0	0	1,476
Wholesale/Clinic/Hypodermic/3PL Received	1,014	757	0	0	1,771
Wholesale/Clinic/Hypodermic/3PL Responded	803	563	0	0	1,366
Automated Drug Delivery Systems Received	180	252	0	0	432
Automated Drug Delivery Systems Responded	112	197	0	0	309
Pharmacist-in-Charge Received	686	679	0	0	1,365
Pharmacist-in-Charge Responded	542	413	0	0	955
Change of Permit Received	1,285	1,090	0	0	2,375
Change of Permit Responded	891	746	0	0	1,637
Renewals Received	1,986	2,135	0	0	4,121
Renewals Responded	1,727	1,869	0	0	3,596

Telephone Calls Received	July - Sept	Oct-Dec	Jan-Mar	Apr-Jun	Total FYTD
Designated Representative	62	12	0	0	74
Advanced Practice Pharmacist	16	9	0	0	25
Pharmacist/Intern	991	980	0	0	1,971
Pharmacy	480	477	0	0	957
Sterile Compounding/Outsourcing	121	77	0	0	198
Wholesale/Clinic/Hypodermic/3PL	232	166	0	0	398
Automated Drug Delivery Systems	40	103	0	0	143
Pharmacist-in-Charge	117	60	0	0	177
Change of Permit	118	107	0	0	225
Renewals	853	985	0	0	1,838
Reception	17,184	17,940	0	0	35,124

UPDATE LICENSING RECORDS

Change of Pharmacist-in-Charge	July - Sept	Oct-Dec	Jan-Mar	Apr-Jun	Total FYTD
Received	441	546	0	0	987
Processed	469	489	0	0	958
Approved	457	471	0	0	928
Pending (Data reflects number of pending at the end of the quarter.)	175	250			n/a
Change of Designated Representative-in-Charge	July - Sept	Oct-Dec	Jan-Mar	Apr-Jun	Total FYTD
Received	28	38	0	0	66
Processed	24	31	0	0	55
Approved	23	24	0	0	47
Pending (Data reflects number of pending at the end of the quarter.)	175	67			n/a
Change of Responsible Manager	July - Sept	Oct-Dec	Jan-Mar	Apr-Jun	Total FYTD
Received	6	2	0	0	8
Processed	5	2	0	0	7
Approved	5	2	0	0	7
Pending (Data reflects number of pending at the end of the quarter.)	2	2			n/a
Change of Professional Director	July - Sept	Oct-Dec	Jan-Mar	Apr-Jun	Total FYTD
Received	24	27	0	0	51
Processed	31	19	0	0	50
Approved	17	12	0	0	29
Pending (Data reflects number of pending at the end of the quarter.)	47	61			n/a
Change of Permits	July - Sept	Oct-Dec	Jan-Mar	Apr-Jun	Total FYTD
Received	238	267	0	0	505
Processed	268	332	0	0	600
Approved	178	475	0	0	653
Pending (Data reflects number of pending at the end of the quarter.)	1,799	1,611			n/a
Clinic Co-Location	July - Sept	Oct-Dec	Jan-Mar	Apr-Jun	Total FYTD
Received	0	0	0	0	0
Processed	0	0	0	0	0
Approved	0	0	0	0	0
Pending (Data reflects number of pending at the end of the quarter.)	0	0			n/a
Discontinuance of Business	July - Sept	Oct-Dec	Jan-Mar	Apr-Jun	Total FYTD
Received	92	101	0	0	193
Processed	92	59	0	0	151
Approved	79	38	0	0	117
Pending (Data reflects number of pending at the end of the quarter.)	237	294			n/a
Intern Pharmacist Extensions	July - Sept	Oct-Dec	Jan-Mar	Apr-Jun	Total FYTD
Received	31	37	0	0	68
Processed	25	33	0	0	58
Completed	13	27	0	0	40
Pending (Data reflects number of pending at the end of the quarter.)	28	40			n/a
Requests Approved	July - Sept	Oct-Dec	Jan-Mar	Apr-Jun	Total FYTD
Address/Name Changes	3,351	3,009	0	0	6,360
Off-site Storage	94	14	0	0	108
Transfer of Intern Hours	5	10	0	0	15
License Verification	514	453	0	0	967

DISCONTINUED OF BUSINESS

discontinued by date of closure

Site Licenses	July - Sept	Oct-Dec	Jan-Mar	Apr-Jun	Total FYTD
Automated Drug Delivery System (ADD)	12	31	0	0	43
Automated Drug Delivery System EMS (ADE)	0	0	0	0	0
Automated Patient Dispensing System 340B Clinic (ADC)	0	1	0	0	1
Centralized Hospital Packaging Government Owned (CHE)	0	0	0	0	0
Centralized Hospital Packaging (CHP)	0	0	0	0	0
Clinics (CLN)	2	3	0	0	5
Clinics Government Owned (CLE)	1	2	0	0	3
Drug Room (DRM)	0	1	0	0	1
Drug Room Government Owned (DRE)	0	0	0	0	0
Hospitals (HSP)	2	1	0	0	3
Hospitals Government Owned (HPE)	1	1	0	0	2
Hospital Satellite Sterile Compounding (SCP)	0	0	0	0	0
Hospital Satellite Sterile Compounding Government Owned (SCE)	0	0	0	0	0
Hypodermic Needle and Syringes (HYP)	2	0	0	0	2
Correctional Pharmacy (LCF)	0	0	0	0	0
Outsourcing Facility (OSF)	0	0	0	0	0
Outsourcing Facility Nonresident (NSF)	0	0	0	0	0
Pharmacy (PHY)	20	30	0	0	50
Pharmacy (PHY) Chain	10	7	0	0	17
Pharmacy Government Owned (PHE)	0	0	0	0	0
Remote Dispensing Pharmacy (PHR)	0	1	0	0	1
Pharmacy Nonresident (NRP)	4	4	0	0	8
Sterile Compounding (LSC)	4	10	0	0	14
Sterile Compounding Government Owned (LSE)	1	0	0	0	1
Sterile Compounding Nonresident (NSC)	0	3	0	0	3
Surplus Medication Collection Distribution Intermediary (SME)	0	0	0	0	0
Third-Party Logistics Providers (TPL)	0	2	0	0	2
Third-Party Logistics Providers Nonresident (NPL)	2	0	0	0	2
Veterinary Food-Animal Drug Retailer (VET)	0	0	0	0	0
Wholesalers (WLS)	3	3	0	0	6
Wholesalers Government Owned (WLE)	0	0	0	0	0
Wholesalers Nonresident (OSD)	4	5	0	0	9
Total	68	105	0	0	173

LICENSES RENEWED

Individual Licenses Renewed	July - Sept	Oct-Dec	Jan-Mar	Apr-Jun	Total FYTD
Designated Representatives (EXC)	495	632	0	0	1,127
Designated Representatives Vet (EXV)	12	7	0	0	19
Designated Representatives-3PL (DRL)	82	62	0	0	144
Designated Representatives-Reverse Distributor (DRR)	0	0	0	0	0
Designated Paramedic (DPM)	0	0	0	0	0
Pharmacist (RPH)	4,846	5,254	0	0	10,100
Advanced Practice Pharmacist (APH)	99	112	0	0	211
Pharmacy Technician (TCH)	6,975	7,640	0	0	14,615
Total	12,509	13,707	0	0	26,216

Site Licenses Renewed	July - Sept	Oct-Dec	Jan-Mar	Apr-Jun	Total FYTD
Automated Drug Delivery System (ADD)	21	645	0	0	666
Automated Drug Delivery System EMS (ADE)	0	0	0	0	0
Automated Patient Dispensing System 340B Clinic (ADC)	0	0	0	0	0
Centralized Hospital Packaging Government Owned (CHE)	2	0	0	0	2
Centralized Hospital Packaging (CHP)	3	1	0	0	4
Clinics (CLN)	400	231	0	0	631
Clinics Government Owned (CLE)	116	800	0	0	916
Drug Room (DRM)	3	4	0	0	7
Drug Room Government Owned (DRE)	5	5	0	0	10
Hospitals (HSP)	53	142	0	0	195
Hospitals Government Owned (HPE)	44	11	0	0	55
Hospital Satellite Sterile Compounding (SCP)	2	1	0	0	3
Hospital Satellite Sterile Compounding Government Owned (SCE)	0	0	0	0	0
Hypodermic Needle and Syringes (HYP)	59	52	0	0	111
Correctional Pharmacy (LCF)	2	57	0	0	59
Outsourcing Facility (OSF)	1	2	0	0	3
Outsourcing Facility Nonresident (NSF)	2	6	0	0	8
Pharmacy (PHY)	1,027	1,994	0	0	3,021
Pharmacy Government Owned (PHE)	59	50	0	0	109
Remote Dispensing Pharmacy (PHR)	1	0	0	0	1
Pharmacy Nonresident (NRP)	69	164	0	0	233
Sterile Compounding (LSC)	139	244	0	0	383
Sterile Compounding Government Owned (LSE)	69	5	0	0	74
Sterile Compounding Nonresident (NSC)	10	18	0	0	28
Surplus Medication Collection Distribution Intermediary (SME)	1	0	0	0	1
Third-Party Logistics Providers (TPL)	4	5	0	0	9
Third-Party Logistics Providers Nonresident (NPL)	20	24	0	0	44
Veterinary Food-Animal Drug Retailer (VET)	3	5	0	0	8
Wholesalers (WLS)	123	100	0	0	223
Wholesalers Government Owned (WLE)	5	5	0	0	10
Wholesalers Nonresident (OSD)	174	173	0	0	347
Total	2,417	4,744	0	0	7,161

CURRENT LICENSES - Data reflects number of licenses at the end of the quarter.

Individual Licenses	July - Sept	Oct-Dec	Jan-Mar	Apr-Jun
Designated Representatives (EXC)	2,849	2,826		
Designated Representatives Vet (EXV)	67	60		
Designated Representatives-3PL (DRL)	351	358		
Designated Representatives-Reverse Distributor (DRR)	4	5		
Designated Paramedic (DPM)	3	3		
Intern Pharmacist (INT)	7,039	6,492		
Pharmacist (RPH)	48,587	48,788		
Advanced Practice Pharmacist (APH)	851	896		
Pharmacy Technician (TCH)	68,637	68,350		
Total	128,388	127,778	0	0

Site Licenses	July - Sept	Oct-Dec	Jan-Mar	Apr-Jun
Automated Drug Delivery System (ADD)	930	942		
Automated Drug Delivery System EMS (ADE)	1	1		
Automated Patient Dispensing System 340B Clinic (ADC)	1	0		
Centralized Hospital Packaging Government Owned (CHE)	2	2		
Centralized Hospital Packaging (CHP)	8	8		
Clinics (CLN)	1,311	1,317		
Clinics Government Owned (CLE)	898	907		
Drug Room (DRM)	22	22		
Drug Room Government Owned (DRE)	10	10		
Hospitals (HSP)	392	401		
Hospitals Government Owned (HPE)	79	79		
Hospital Satellite Sterile Compounding (SCP)	4	4		
Hospital Satellite Sterile Compounding Government Owned (SCE)	2	2		
Hypodermic Needle and Syringes (HYP)	300	300		
Correctional Pharmacy (LCF)	61	61		
Outsourcing Facility (OSF)	4	4		
Outsourcing Facility Nonresident (NSF)	25	25		
Pharmacy (PHY)	6,378	6,387		
Pharmacy Government Owned (PHE)	136	136		
Remote Dispensing Pharmacy (PHR)	3	2		
Pharmacy Nonresident (NRP)	589	583		
Sterile Compounding (LSC)	744	742		
Sterile Compounding Government Owned (LSE)	112	112		
Sterile Compounding Nonresident (NSC)	68	66		
Surplus Medication Collection Distribution Intermediary (SME)	1	1		
Third-Party Logistics Providers (TPL)	32	33		
Third-Party Logistics Providers Nonresident (NPL)	84	92		
Veterinary Food-Animal Drug Retailer (VET)	20	20		
Wholesalers (WLS)	530	530		
Wholesalers Government Owned (WLE)	14	14		
Wholesalers Nonresident (OSD)	807	809		
Total	13,568	13,612	0	0
Total Population of Licenses	141,956	141,390	0	0